

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

To

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C.20231
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

Date of mailing (day/month/year)

08 March 2000 (08.03.00)

International application No.

PCT/US99/13411

Applicant's or agent's file reference

KIMCHI2APCT

International filing date (day/month/year)

15 June 1999 (15.06.99)

Priority date (day/month/year)

15 June 1998 (15.06.98)

Applicant

KIMCHI, Adi

1. The designated Office is hereby notified of its election made:



in the demand filed with the International Preliminary Examining Authority on:

14 January 2000 (14.01.00)



in a notice effecting later election filed with the International Bureau on:

2. The election



was



was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

F. Baechler

Telephone No.: (41-22) 338.83.38

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT


(PCT Article 36 and Rule 70)

Applicant's or agent's file reference KIMCH12APCT	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US99/13411	International filing date (day/month/year) 15 JUNE 1999	Priority date (day/month/year) 15 JUNE 1998
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant YEDA RESEARCH AND DEVELOPMENT COMPANY LTD.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 3 sheets.
☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).
 These annexes consist of a total of 0 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☒ Non-establishment of report with regard to novelty, inventive step or industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 14 JANUARY 2000	Date of completion of this report 21 AUGUST 2000
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer MARYAM MONSHIPOURI  Telephone No. (703) 308-0196
Facsimile No. (703) 305-3230	

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/13411

I. Basis of the report**1. With regard to the elements of the international application:***☒ the international application as originally filed☒ the description:

pages 1-35 , as originally filed
pages NONE , filed with the demand
pages NONE , filed with the letter of _____

☒ the claims:

pages 36-39 , as originally filed
pages NONE , as amended (together with any statement) under Article 19
pages NONE , filed with the demand
pages NONE , filed with the letter of _____

☒ the drawings:

pages 1-10 , as originally filed
pages NONE , filed with the demand
pages NONE , filed with the letter of _____

☒ the sequence listing part of the description:

pages 1-12 , as originally filed
pages NONE , filed with the demand
pages NONE , filed with the letter of _____

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

- ☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).
☐ the language of publication of the international application (under Rule 48.3(b)).
☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:☒ contained in the international application in printed form.☒ filed together with the international application in computer readable form.☐ furnished subsequently to this Authority in written form.☐ furnished subsequently to this Authority in computer readable form.☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.**4. ☒ The amendments have resulted in the cancellation of:**☒ the description, pages NONE☒ the claims, Nos. NONE☒ the drawings, sheets/fig NONE**5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).****

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

**Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/13411

III Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non obvious), or to be industrially applicable have not been and will not be examined in respect of:

☐ the entire international application.

☒ claims Nos. 21-22, 25-26

because:

☐ the said international application, or the said claim Nos. _ relate to the following subject matter which does not require international preliminary examination (*specify*).

☐ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. _ are so unclear that no meaningful opinion could be formed (*specify*).

☐ the claims, or said claims Nos. _ are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for said claims Nos. 21-22, 25-26.

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/13411

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**1. statement****Novelty (N)**Claims 3-7, 10, 13-17 YESClaims 1-2, 8-9, 11-12, 18-20, 23-24 NO**Inventive Step (IS)**Claims 3-7, 10, 13-17 YESClaims 1-2, 8-9, 11-12, 18-20, 23-24 NO**Industrial Applicability (IA)**Claims 1-20, 23-24 YESClaims NONE NO**2. citations and explanations (Rule 70.7)**

Claims 1-2, 8-9, 11-12, and 18-20 lack novelty under PCT Article 33(2) as being anticipated by Yeda Research and Development Co. (WO 95/10630, 20 April 1995). Kimchi et al. teach a nucleic acid molecule encoding DAP kinase comprising a region which has at least 85% amino acid identity to the kinase domain of SEQ ID NO:2 and is capable of inducing cell death and can be considered to be a fragment of DRP-1 of this invention, anticipating claim 1 (and its dependent claim 2) as well as claim 8 (and its dependent claims 9, 10 and 18-20). Further, their nucleotide sequence hybridizes to bases 96-886 of SEQ ID NO:1 under stringent hybridization conditions, anticipating claims 11-12 (it should be noted that applicant did not define the hybridization and wash conditions corresponding to "moderately" or "high" stringent conditions and therefore those were assumed to basically refer to "stringent hybridization conditions").

Claims 1-2, 8-9, 11-12, 18-20 and 23-24 lack novelty under PCT Article 33(2) as being anticipated by Deiss et al. (Genes and Development, 9, 15-30, 1995). Deiss et al. teach a nucleic acid sequence encoding DAP kinase comprising a region which has at least 85% amino acid identity to the kinase domain of SEQ ID NO:2 and is capable of inducing cell death and can be considered to be a fragment of DRP-1 of this invention anticipating claim 1 (and its dependent claim 2) as well as claim 8 (and its dependent claims 9, 10 and 18-20). Their nucleic acid sequence comprises a portion of the mRNA corresponding to SEQ ID NO:2 and is capable of hybridizing to said RNA to prevent its translation, anticipating claim 23. Further, their nucleotide sequence hybridizes to bases 96-886 of SEQ ID NO:1 under stringent hybridization conditions, anticipating claims 11-12 (it should be noted that applicant did not define the hybridization and wash conditions corresponding to "moderately" or "high" stringent conditions and therefore those were assumed to basically refer to "stringent hybridization conditions"). Deiss et al. specifically teach a method of neutralizing mRNA molecule, comprising contacting a single strand RNA fragment (Continued on Supplemental Sheet.)

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/13411

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

CLASSIFICATION:

The International Patent Classification (IPC) and/or the National classification are as listed below:

IPC(7): C12N 9/12, 1/20, 5/00, 15/00; C12Q 1/68; C07H 21/04; A61K 38/51 and US Cl.: 435/194, 320.2, 325, 252.3, 6; 424/94.5; 536/23.2

V. 2. REASONED STATEMENTS - CITATIONS AND EXPLANATIONS (Continued):

corresponding to a portion of SEQ ID NO:2 by hybridizing and preventing its translation into complete SEQ ID NO:2. They in fact used this approach to identify SEQ ID NO:2 (see page 19), anticipating claim 24.

Claims 3-7, 10 and 13-17 meet the novelty and inventive requirements of PCT Articles 33(2) and 33(3). This is because these claims are directed to nucleic acid sequences encoding DRP-1 and their expression products of specific composition. Said specifically claimed nucleic acid/amino acid sequences are free of prior art. Further the prior art does not suggest preparing such specific DNA/amino acid sequences. Hence said sequences are also non-obvious.

Claims 1-20 and 23-24 meet the criteria set out in PCT Article 33(4), for industrial applicability.

NEW CITATIONS

NONE

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/12, 1/20, 5/00, 15/00, C12Q 1/68, C07H 21/04, A61K 38/51		A1	(11) International Publication Number: WO 99/66030
			(43) International Publication Date: 23 December 1999 (23.12.99)
(21) International Application Number: PCT/US99/13411 (22) International Filing Date: 15 June 1999 (15.06.99) (30) Priority Data: 60/089,294 15 June 1998 (15.06.98) US (71) Applicant (for all designated States except US): YEDA RESEARCH AND DEVELOPMENT COMPANY LTD. [IL/IL]; P.O. Box 96, 76100 Rehovot (IL). (71) Applicant (for SD only): MCINNIS, Patricia, A. [US/US]; Apartment #203, 2325 42nd Street N.W., Washington, DC 20007 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): KIMCHI, Adi [IL/IL]; 38 Hashalom Street, 43561 Raanana (IL). (74) Agent: BROWDY, Roger, L.; Browdy and Neimark, P.L.L.C., Suite 300, 419 Seventh Street N.W., Washington, DC 20004 (US).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: DAP-KINASE RELATED PROTEIN			
(57) Abstract A new protein, which is a novel homologue of DAP-kinase, has been isolated. This novel calmodulin-dependent kinase is a cell death-promoting protein functioning in the biochemical pathway which involves DAP (death-associated protein)-kinase (e.g., forming a cascade of sequential kinases, one directly activating the other). Alternatively, the two kinases may operate to promote cell death in parallel pathways.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DAP-KINASE RELATED PROTEIN**CROSS-REFERENCE TO RELATED APPLICATIONS**

5 The present application is related to and hereby incorporates by reference the entire contents of each of application serial no. 08/810,712, filed March 3, 1997, and application serial no. 08/631,097, filed April 12, 1996, the latter of which has been published as PCT publication WO 95/10630 on April 20, 1995. The present application also claims priority from provisional application 60/089,294, filed June 15, 1998, the entire contents of which are hereby
10 incorporated by reference.

BACKGROUND OF THE INVENTIONField of the Invention

 The present invention is directed to a DAP-kinase related protein.

15

Description of the Related Art

 One of the factors which determines the proliferation state of cells is the balance between the growth-promoting effects of proto-oncogenes and the growth-constraining effects of tumor-suppressor genes. One mechanism by which
20 tumor-suppressor genes exert their growth-constraining effect is by inducing a cell to undergo a physiological type of death. Such a controlled cell death is evident in a multitude of physiological conditions including metamorphosis, synaptogenesis of neurons, death of lymphocytes during receptor repertoire selection, and controlled homeostasis in the bone marrow and other proliferative tissues, etc. This cell death is regulated by the interaction of
25 the cell with other cells or with cell products, for example through the activity of suitable cytokines.

 Growth-inhibiting cytokines have a double effect on the target cell. They can either inhibit the proliferation of the cell and/or give rise to cell death. To date, blockage or activation of expression of known tumor-suppressor genes was shown to counteract or
30 enhance, respectively, cytokines inhibition of cells growth (Kimchi, 1992) but did not have any effect on the death- promoting action of cytokines. For example, the growth inhibitory response to cytokines, such as TGF- β , was markedly reduced by the inactivation of the Rb gene, or the response to IL-6 was enhanced by introducing activated p53 genes (Pietenpol et al., 1990; Levy et al, 1993).

Apoptosis is a genetically controlled cell death process which is important in various developmental stages, as well as for cell maintenance and tissue homeostasis (Jacobson et al., 1997). During the last few years, many of the key players in this process have been identified, including receptors, adaptor proteins, proteases, and other positive and negative regulators (Green et al., 1998; White, 1996). One of the positive mediators of apoptosis, which has been cloned by the present inventors, is DAP-kinase (Deiss et al., 1995). This protein was discovered by a functional approach to gene cloning, based on transfections of mammalian cells with anti-sense cDNA libraries and subsequent isolation of death-protective cDNA fragments (Deiss et al., 1995; Deiss et al., 1991; Kimchi, 1998; Kissil et al., 1998; Levy-Strumpf et al., 1998). The anti-sense cDNA of DAP-kinase protected HeLa cells from interferon-gamma-induced cell death, and this property served as the basis for its selection.

DAP-kinase is a calcium/calmodulin-regulated 160 kDa serine/threonine protein kinase associated with actin microfilaments (Deiss et al., 1995; Cohen et al., 1997). Its structure contains at least two additional domains that might mediate interactions with other proteins: ankyrin repeats, and a typical death domain located at the C-terminal part of the protein (Deiss et al., 1995; Cohen et al., 1997). Overexpression of DAP-kinase in various cell lines results in cell death, and this death-promoting effect of DAP-kinase depends on at least three features: the catalytic activity, presence of the death domain, and the correct intracellular localization (Cohen et al., 1997; Cohen et al., 1999). Several independent lines of evidence proved that DAP-kinase is involved in apoptosis triggered by different external signals, including interferon- γ , TNF- α , activated Fas receptors, and detachment of cells from the extracellular matrix (Deiss et al., 1995; Cohen et al., 1997; Cohen et al., 1999; Inbal et al., 1997). A tumor suppressive function was recently attributed to the DAP-Kinase, coupling the control of apoptosis to metastasis (Inbal et al., 1997).

So far, only a few serine/threonine kinases were implicated in the regulation of programmed cell death, either as death-promoting and death-protecting proteins (Anderson, 1997; Bokoch, 1998). One such candidate is the JNK/SAPK (Basu et al., 1998). In one example, it was shown to mediate apoptosis induced by detachment from extracellular matrix (named anoikis) (Cardone et al., 1997). In this system, the JNK pathway is activated by MEKK-1, whose kinase activity is stimulated by caspase cleavage (Cardone et al., 1997). JNK

may antagonize BCL-2 anti-apoptotic effects by phosphorylation (Park et al., 1997; Maundrell et al., 1997).

Another serine/threonine kinase is RIP, which like DAP-Kinase also possesses the death domain. RIP was shown to positively mediate apoptosis in cell cultures (Stanger et al., 1995). However, *in vivo* studies in RIP-deficient mice demonstrated its ability to exert anti-apoptotic effects by mediating the TNF- α -induced TNF- β activation (Kelliher et al., 1998). Other RIP members, RIP2 and RIP 3 were also recently identified and shown to possess pro-apoptotic effects (McCarthy et al., 1998; Sun et al., 1998; Yu et al., 1999).

Among the negative regulators of apoptosis is the protein kinase AKT. This protein was shown to phosphorylate BAD and thereby to prevent it from complexing and blocking the anti-apoptotic activity of BCL-X_L (Datta et al., 1997; del Peso et al., 1997). AKT was also recently shown to phosphorylate pro-caspase-9, thus blocking its normal processing (Cardone et al., 1998).

Recently, the isolation and characterization of novel kinase members, homologous in their catalytic domains to DAP-kinase, was reported (Kawai et al., 1998; Kogel et al., 1998; Sanjo et al., 1998). One protein, named ZIP-kinase, was found to be 80% identical to DAP-kinase within the kinase domain, yet it lacks the CaM-regulatory domain and the other domains and motifs characteristic of DAP-kinase. Zip-kinase contains a leucine zipper domain at the C-terminus and is localized to the nucleus (Kawai et al., 1998; Kogel et al., 1998). The activation of ZIP kinase occurs by a different mechanism involving homo-dimerization, mediated by its leucine zipper domain. However, unlike DAP-kinase, ZIP-kinase is a nuclear protein, which instead of being regulated by a calmodulin-binding domain, is activated by homo-dimerization of its leucine-zipper motifs (Kogel et al., 1998). Another two less conserved nuclear proteins, DRAK1 and DRAK2, which are closely related to each other, and which share 50% identity with the kinase domain of DAP-kinase, were also recently characterized. Like ZIP-kinase, the DRAK1 and DRAK2 proteins also lack the CaM-regulatory domain. The overexpression of these two proteins in NIH3T3 cells induces some morphological changes associated with apoptosis, dependent on the functionality of their kinase domain (Sanjo et al., 1998). Together these kinases form a novel subfamily of serine/threonine kinases, as is evident from multiple sequence and phylogenetic analysis (Inbal et al., 1999).

Ectopic expression of the three wild type kinases, but not their catalytically inactive mutants, induced morphological changes characteristic of apoptosis (Kawai et al., 1998; Sanjo et al., 1998). Yet, in the case of ZIP-Kinase, these results are still controversial (Kogel et al., 1998).

5 Citation of any document herein is not intended as an admission that such document is pertinent prior art, or considered material to the patentability of any claim of the present application. Any statement as to the content or a date of any document is based on information available to the applicant at the time of filing and does not constitute an admission as to the correctness of such a statement

SUMMARY OF THE INVENTION

10 A new protein, DAP-Kinase-related 1 protein (DRP-1), which is a novel homologue of DAP-kinase, has been isolated. This novel calmodulin-dependent kinase is a 42kDa serine/threonine kinase which shows a high degree of homology to DAP-kinase both in
15 its catalytic domain and its calmodulin-regulatory region. The catalytic domain of DRP-1 is also homologous to recently identified ZIP-kinase and, to a lesser extent, to the catalytic domains of DRAK1/2.

DRP-1 is localized to the cytoplasm as shown by immunostaining and cellular fractionation assays. *In vitro* kinase assays indicate that wild type DRP-1, but not a kinase
20 inactive mutant, undergoes autophosphorylation and phosphorylates an external substrate in a Ca²⁺/CaM-dependent manner. Ectopically expressed DRP-1 is able to induce apoptosis in various types of cells; with this killing being dependent on its kinase activity. The dominant negative form of DAP-Kinase (DAPk DD) is a potent blocker of apoptosis induced DRP-1. Thus, DRP-1 may be a death-promoting protein functioning in the biochemical pathway which
25 involves DAP (death-associated protein)-kinase (e.g., forming a cascade of sequential kinases, one directly activating the other). Alternatively, the two kinases may operate to promote cell death in parallel pathways.

The present invention provides for a DRP-1 protein and functional homologues thereof having at least 85% sequence identity to the DRP-1 sequence of SEQ ID NO:2. Also
30 provided is a fragment of DRP-1, which either is capable of inducing cell death or lacks such capability but instead is capable of inhibiting the activity of DRP-1 or a functional homologue

thereof to induce cell death, and a homologous fragment which has at least 85% sequence identity thereto and which has the same properties.

The present invention further provides an isolated DNA molecule encoding for such DRP-1 protein, functional homologues thereof, or fragments thereof. Also included within the scope of the present invention are isolated DNA molecules which hybridize to the nucleotide sequence encoding DRP-1 protein under moderately or highly stringent conditions and encode a calmodulin-dependent serine/threonine kinase having the property of being capable of inducing cell death.

Other further aspects of the present invention include a composition comprising the DRP-1 protein, functional homologues and fragments thereof, and an antibody which specifically recognizes DRP-1 but does not cross-react with DAP kinase or ZIP kinase.

Yet another aspect of the present invention is directed to a single stranded RNA molecule complementary to at least a portion of the mRNA encoding the DRP-1 protein of SEQ ID NO:2. This single stranded antisense RNA molecule can be used in a method of neutralizing DRP-1 mRNA by hybridizing to the DRP-1 mRNA to prevent its translation into DRP-1 protein.

The present invention also provides a method for screening individuals for predisposition to cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide (SEQ ID NO:1) and amino acid (SEQ ID NO:2) sequence of the DAP-kinase homologue, DRP-1. The initiation (ATG) and stop (TAA) codons are boxed. The polyadenylation signal (ATTAAA) is underlined. The kinase domain and the calmodulin regulatory regions are in bold or underlined by a dash, respectively.

Figures 2A-2B show the multiple sequence alignment of the serine/threonine kinase domains (Fig. 2A) of the DAP-kinase-related proteins, DAP-kinase (SEQ ID NO:3), ZIP-kinase (SEQ ID NO:4), DRP-1 (corresponding to residues 13-275 of SEQ ID NO:2), DRAK1 (SEQ ID NO:5) and DRAK 2 (SEQ ID NO:6), conducted according to Hanks and Quinn (1991) with identical amino acids boxed and homologous amino acids shown with gray shading, and the multiple sequence alignment of the calmodulin regulatory regions (Fig. 2B) of DAP-kinase (SEQ ID NO:7), DRP-1 (corresponding to residues 292 to 320 of SEQ ID NO:2), smMLCK (SEQ ID NO:8), CaMKIIa (SEQ ID NO:9), CaMKI (SEQ ID NO:10),

CaMKIV (SEQ ID NO:11), and ZIP-Kinase (SEQ ID NO:12) conducted manually, keeping the conserved (boxed) regions aligned to each other. The corresponding region of ZIP-Kinase which does not contain homology to DAP-Kinase and ZIP-Kinase CAM-regulatory regions is given at the bottom of Fig. 2B.

5 Figure 3A shows Northern blot analysis of polyA+RNA extracted from various cell lines for mRNA expression of DRP-1, Figure 3B show Western blot analysis of *in vitro* transcription and translation of DRP-1, and Figure 3C shows protein expression of DRP-1 in HeLA cells on an immunoblot.

10 Figures 4A and 4B show control COS-7 cells and cellular localization of DRP-1 in COS-7 cells, respectively, and Figure 4C shows a Western blot of fractions from a detergent extraction of COS-7 cells transfected with a pCDNA3 vector expressing either FLAG-tagged DRP-1 or DAP-Kinase.

 Figure 5A shows *in vitro* kinase activity of DRP-1 and Figure 5B shows a Western blot of DRP-1 proteins.

15 Figures 6A-6B show fluorescent microscope images of 293 cells transfected by pCDNA3-luciferase as a negative control (Fig. 6A), by pCDNA3- Δ CaM DAP-Kinase as positive control (Fig. 6B), by pCDNA -DRP-1 (Fig. 6C), and by pCDNA3-K42A DRP-1 (Fig. 6D). Apoptotic cells are indicated by arrows.

20 Figure 7 shows the scores of apoptotic cells in a graph of the percentage of apoptotic cells resulting from the transfections of Figs. 6A-6D.

 Figures 8A and 8B show DRP-1 protein expression in 293 transfected cells in immunoblots to anti-FLAG antibodies (Fig. 8A) and anti-vinculin antibodies (Fig. 8B).

25 Figure 9A shows that DAP kinase death domain protects from DRP-1 induced apoptosis, and Figure 9B shows an immunoblot of DRP-1 protein expression in 293 transfected cells.

 Figure 10A shows a schematic representation of a series of generated deletion mutant, and Figure 10B shows an immunoblot containing extracts of 293 cells transiently transfected with GFP and the series of deletion mutants, (DRP-1 fragments, cloned in pCDNA3, and tagged with HA epitope at the C-terminus), as in Figs. 8A and 8B are probed
30 with anti-HA antibodies for DRP-1 detection and anti-vinculin antibodies to quantitate the loaded protein amounts. In Figure 10B, pCDNA3²-luciferase is the negative control.

Figure 11A shows fluorescent microscope images of the transiently transfected cells of Fig. 10B, and Figure 11B shows a graph of the score in percent apoptotic cells in Fig. 11A resulting from co-transfections of 293 cells with 1-2 μ g HA-tagged wild type DRP-1 or various deletion mutants of DRP-1 after 24 hours (average S.D. calculated from triplicates of cells each).

Figures 12A and 12B show by Western analysis that the C-terminal part of DRP-1 is required for its homo-dimerization. In Figure 12A, wild type DRP-1 is shown to undergo specific homo-dimerization. The lanes correspond to the following co-transfections of DRP-1 constructs and 20 μ g of RFX1- Δ SmaI constructs/9mm plate): (1) DRP-1-FLAG+RFX1- Δ SmaI-HA (control to rule-out nonspecific attachment of DRP-1 to HA beads or an irrelevant gene). (2) RFX- Δ SmaI-FLAG+DRP-1-HA (control to rule out nonspecific attachment of DRP-1 to FLAG beads or to an irrelevant gene). (3) DRP-1-FLAG+DRP-1-FLAG. Both IP directions and their Western blottings are shown. In Figure 12B, truncation of the final 40 amino acids of DRP-1 is shown to abolish its homo-dimerization. The lanes correspond to the following co-transfections (5 μ g of each construct/90mm plate): (1) DRP-1-FLAG+DRP-1-HA (2) DRP-1-FLAG+DRP-1- Δ 40-HA (3) DRP-1-FLAG+DRP-1- Δ 73-HA (4) DRP-1-FLAG+DRP-1- Δ 85-HA. The lower panel quantitate the immunoprecipitation efficiency of DRP-1-FLAG by the anti-FLAG antibodies.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the discovery by the present inventor of a novel serine/threonine kinase with remarkable homology to the catalytic and CaM-regulatory domains of DAP-kinase. This kinase, named DAP-kinase-related protein 1 (DRP-1), is a 42kDa cytoplasmic protein which exhibits minor associations with insoluble matrix elements. The nucleotide (SEQ ID NO:1) and amino acid (SEQ ID NO:2) sequences of this DRP-1 protein are shown in Fig 1. It is composed of 1742 nucleotides. The predicted initiation and stop codons are boxed, and the polyadenylation signal is underlined. The protein kinase domain is shown in bold letters and corresponds to amino acid residues 13 to 275 of SEQ ID NO:2. This protein displays 80% identity with the catalytic domain of DAP-kinase. The calmodulin-regulatory region is underlined with a dashed line; this region displays high homology to the corresponding region in DAP-kinase. The remainder of the C-terminal short part of the protein (the last 40 amino acid residues corresponding to residues 321 to 360 of

SEQ ID NO:2) differs completely from DAP-kinase. Thus, DAP-kinase-related I protein does not carry all of the other motifs and protein modules characteristic of DAP-kinase. The mRNA expression levels transcribed from this gene are low.

Another protein, ZIP-kinase, which by virtue of its sequence homology to the kinase domain of DAP-Kinase, is also a member of the DAP-Kinase-related proteins subfamily, was recently identified (Kawai et al., 1998; Kogel et al., 1998). Unlike DAP-Kinase and DRP-1, ZIP-kinase is a nuclear protein, which instead of being regulated by a calmodulin-binding domain, is activated only by homo-dimerization via its leucine-zipper motifs (Kawai et al., 1998). To this group of kinases, another two less homologous nuclear proteins, DRAK1 and DRAK2, were recently added (Sanjo et al., 1998). Together they form a novel subfamily of serine/threonine kinases, as is evident from multiple sequence and phylogenetic analyses (Inbal et al., 1999). A multiple sequence alignment of the kinase domain of these serine/threonine kinases is shown in Fig. 2A.

To check the cellular functions of DRP-1, the laboratory of the present inventor overexpressed wild type DRP-1 in various cell lines and found that it induced apoptosis as measured by a few parameters. Unlike the wild type DRP-1, a kinase inactive mutant of DRP-1 (DRP-1 K42A), did not induce apoptosis, although it was expressed at a similar level in the transfected cells. *In vitro* kinase assays confirmed that DRP-1 K42A is indeed unable to phosphorylate MLC or under autophosphorylation. Also, a truncated form of DRP-1 which lacks the CaM-regulatory region, could be shown to induce very high levels of apoptosis, in a similar way to the analogous truncation of the CaM-regulatory region of DAP-Kinase (Δ CaM; Cohen et al., 1997). Such dependence on the catalytic activity for the apoptotic function is apparent also in the other members of DAP-kinase-related proteins (Kawai et al., 1998; Sanjo et al., 1998).

In a deletion mutant study, which is also presented in the Example herein, the existence of a positive element responsible for apoptotic induction which is located at the C-terminal part of DRP-1 is confirmed. This C-terminal tail of DRP-1 is also essential for its dimerization. Thus, self-dimerization is a requirement for the functionality of this kinase in apoptotic assays, although this property can be overridden by a further deletion of the CaM-regulatory region. Like DRP-1, ZIP-kinase-induced cell death is also controlled by its ability to undergo homo-dimerization via the C-terminal leucine zipper domain (Kawai et al., 1998). Three point mutations in the leucine zipper domain abolished the homo-dimerization as well as

the ability of ZIP-kinase to undergo autophosphorylation *in vitro* and significantly reduced its ability to induce cell death of NIH 3T3 cells. It seems reasonable to assume that activation of these kinases is achieved by homo-dimerization followed by trans-phosphorylation events.

The high homology in the kinase domains of DAP-kinase and DRP-1, and the
5 finding that they are both localized to the cytoplasm (either in soluble or insoluble forms), imply that they may share the same or closely related substrates. The phosphorylation sites for these kinases on the substrate may be either different or identical. Thus, these kinases may cooperate to induce apoptosis in the same cell type or, alternatively, function independently in different cell types, tissues or organs, or in response to different stimuli or time windows.
10 Another possibility is that these kinases act sequentially along the same signaling pathway to induce apoptosis.

The present invention thus provides for the polypeptide of DRP-1 and for a calmodulin-dependent serine/threonine kinase homologue having the properties of DRP-1, such as the ability to phosphorylate protein in a calcium/calmodulin dependent manner and the
15 ability to induce programmed cell death or apoptosis, and having at least 85% sequence identity to the amino acid sequence SEQ ID NO:2 of DRP-1. Preferably, the calmodulin-dependent serine/threonine kinase homologue has at least 90% sequence identity, and more preferably, at least 95% sequence identity to SEQ ID NO:2.

The term "sequence identity" as used herein means that the amino acid
20 sequences are compared by alignment according to Hanks and Quinn (1991) with a refinement of low homology regions using the Clustal-X program. Such an amino acid alignment is shown in Figs. 2A and 2B where the identical amino acid residues are presented in boxes (cutoff=50%) and homologous amino acid residues, determined according to the PAM 250 matrix, are presented by gray shading (cutoff=65%).

25 The Clustal-X program referred to in the previous paragraph is the Windows interface for the ClustalW multiple sequence alignment program (Thompson et al., 1994). The Clustal-X program is available over the internet at <ftp://ftp-igbmc.u-strasbg.fr/pub/clustalx/>. Of course, it should be understood that if this link becomes inactive, those of ordinary skill in the art can find versions of this program at other links using standard internet search
30 techniques without undue experimentation. Unless otherwise specified, the most recent version of any program referred herein, as of the effective filing date of the present application, is the one which is used in order to practice the present invention.

If the above method for determining "sequence identity" is considered to be nonenabled for any reason, then one may determine sequence identity by the following technique. The sequences are aligned using Version 9 of the Genetic Computing Group's GDAP (global alignment program), using the default (BLOSUM62) matrix (values -4 to +11) with a gap open penalty of -12 (for the first null of a gap) and a gap extension penalty of -4 (per each additional consecutive null in the gap). After alignment, percentage identity is calculated by expressing the number of matches as a percentage of the number of amino acids in the claimed sequence.

In addition to the full length polypeptide of DRP-1 or a functional homologue thereto with at least 85% sequence identity, the present invention also provides for a fragment of the DRP-1 protein of SEQ ID NO:2 which either maintains the ability to induce cell death or lacks this ability but instead is capable of inhibiting the cell killing ability of DRP-1 protein or its functional homologue described above. It was unexpectedly discovered by the present inventor that the 40 amino acid C-terminal tail (residues 321 to 360 of SEQ ID NO:2) is critical to induction of cell death. As the action of DRP-1 is dependent on dimerization, the 40 amino acid tail, by itself, can inhibit the ability of DRP-1 to induce cell death by interfering with and preventing DRP-1 from dimerizing. Furthermore, it was also unexpectedly discovered that the catalytic domain, by itself (without the calmodulin regulatory domain and the 40 amino acid C-terminal tail, e.g., amino acid residues 13 to 275 of SEQ ID NO:2), is super-killing. One of ordinary skill in the art can readily obtain fragments of the full length sequence of the present invention using N-terminal amino peptidases or C-terminal carboxypeptidases. Each fragment can then be readily tested to see if it possesses one of the two functions described herein for such fragments, without undue experimentation.

Besides fragments of DRP-1 having the above-mentioned properties, fragments having an amino acid sequence with at least 85% sequence identity to the above fragments of DRP-1, preferably with at least 90% sequence identity, and more preferably with at least 95% sequence identity, and maintaining the cell death induction or inhibition properties of the original fragment, are also comprehended by the present invention.

Also comprehended by the present invention are chemical derivatives of the DRP-1 and functional homologues and fragments thereof, as defined above, where a "chemical derivative" contains additional chemical moieties not normally part of the DRP-1 amino acid sequence. Covalent modifications of the amino acid sequence are included within

the scope of this invention. Such modifications may be introduced into DRP-1 or fragments thereof by reacting targeted amino acid residues of the peptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues.

Cysteinyl residues most commonly are reacted with alpha-haloacetates (and
5 corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxylmethyl or carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with bromotrifluoroacetone, alpha-bromo- beta-(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl-2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4- nitrophenol, or chloro-7-nitrobenzo-
10 2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylprocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Parabromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0.

15 Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing alpha-amino acid-containing residues include imidoesters, such as methyl picolinimide, pyridoxal phosphate, pyridoxal, chloroborohydride, trinitrobenzenesulfonic acid, O-methylisourea,
20 2,4-pentanedione, and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3- butanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the
25 groups of lysine, as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues *per se* has been studied extensively, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and e-nitro
30 derivatives, respectively.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides (R'N-C-N-R') such as 1-cyclohexyl-3-[2-morpholinyl-(4-ethyl)]

carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl)carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

5 Glutaminyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

The present invention also comprehends an isolated DNA molecule which includes a nucleotide sequence encoding the DRP-1 protein of SEQ ID NO:2, a functional
10 homologue thereof as described above, or a fragment of DRP-1 which either maintains the ability of DRP-1 to induce cell death or lacks this ability but is instead capable of inhibiting the cell killing ability of DRP-1 protein, as defined above. The isolated DNA molecule according to the present invention is also intended to comprehend a DNA molecule which hybridizes under moderately stringent, preferably highly stringent, conditions to the nucleotide sequence
15 encoding DRP-1 (corresponding to nucleotides 62 to 1141 of SEQ ID NO:1) and which encodes a polypeptide which maintains the cell death induction properties of DRP-1. The present invention further comprehends isolated DNA molecules which hybridize under moderately stringent, preferably highly stringent, conditions to a nucleotide sequence which encodes for a fragment of DRP-1 which either maintains the ability of DRP-1 to induce cell
20 death (i.e., nucleotides 98 to 886 of SEQ ID NO:1 encoding the catalytic kinase domain of DRP-1) or lacks the ability but is instead capable of inhibiting the cell killing ability of DRP-1 protein (i.e., nucleotides 1022 to 1141 of SEQ ID NO:1 encoding the 40 amino acid C-terminal tail of DRP-1). Furthermore, polypeptides encoded by any nucleic acid, such as DNA or RNA, which hybridizes to the nucleotide sequence of nucleotides 62 to 141 of SEQ ID
25 NO:1 under moderately stringent or highly stringent conditions are considered to be within the scope of the present invention as long as the encoded polypeptide maintains the ability of DRP-1 to induce cell death.

As used herein, stringency conditions are a function of the temperature used in the hybridization experiment, the molarity of the monovalent cations and the percentage of
30 formamide in the hybridization solution. To determine the degree of stringency involved with any given set of conditions, one first uses the equation of Meinkoth et al. (1984) for

determining the stability of hybrids of 100% identity expressed as melting temperature T_m of the DNA-DNA hybrid:

$$T_m = 81.5^{\circ}\text{C} + 16.6 (\text{Log}M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$$

where M is the molarity of monovalent cations, %GC is the percentage of G and C nucleotides

5 in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. For each 1°C that the T_m is reduced from that calculated for a 100% identity hybrid, the amount of mismatch permitted is increased by about 1%.

Thus, if the T_m used for any given hybridization experiment at the specified salt and formamide concentrations is 10°C below the T_m calculated for a 100% hybrid according to
10 the equation of Meinkoth, hybridization will occur even if there is up to about 10% mismatch.

As used herein, "highly stringent conditions" are those which provide a T_m which is not more than 10°C below the T_m that would exist for a perfect duplex with the target sequence, either as calculated by the above formula or as actually measured.

"Moderately stringent conditions" are those which provide a T_m which is not more than 20°C
15 below the T_m that would exist for a perfect duplex with the target sequence, either as calculated by the above formula or as actually measured. Without limitation, examples of highly stringent (5 - 10°C below the calculated or measured T_m of the hybrid) and moderately stringent (15 - 20°C below the calculated or measured T_m of the hybrid) conditions use a wash solution of 2 X SSC (standard saline citrate) and 0.5% SDS (sodium dodecyl sulfate) at the
20 appropriate temperature below the calculated T_m of the hybrid. The ultimate stringency of the conditions is primarily due to the washing conditions, particularly if the hybridization conditions used are those which allow less stable hybrids to form along with stable hybrids. The wash conditions at higher stringency then remove the less stable hybrids. A common hybridization condition that can be used with the highly stringent to moderately stringent wash
25 conditions described above is hybridization in a solution of 6 X SSC (or 6 X SSPE (standard saline-phosphate-EDTA)), 5 X Denhardt's reagent, 0.5% SDS, 100 $\mu\text{g/ml}$ denatured, fragmented salmon sperm DNA at a temperature approximately 20° to 25°C below the T_m . If mixed probes are used, it is preferable to use tetramethyl ammonium chloride (TMAC) instead of SSC (Ausubel, 1987, 19989).

30 Additional aspects of the present invention are vectors which carry the isolated DNA molecule according to the present invention and a host cell which is transformed with the isolated DNA according to the present invention.

The present invention further provides for antisense RNA complementary to at least a portion of a messenger RNA (mRNA or "sense" RNA) molecule which is the transcription product of the DNA sequence encoding the DRP-1 protein of SEQ ID NO:2. The antisense DRP-1 sequence can be chemically synthesized or it can be expressed in host cells. However, when expressed in host cells, the expressed antisense RNA must be stable (i.e., does not undergo rapid degradation). Moreover, the antisense DRP-1 RNA, will essentially specifically only hybridize to the sense DRP-1 mRNA and form a stable double-stranded RNA molecule that is essentially non-translatable. In other words, the antisense DRP-1 RNA prevents the expressed sense DRP-1 mRNA from being translated into active protein. When expressed in host cells, a vector-borne antisense DRP-1 sequence may contain either the entire DRP-1 gene sequence or merely a portion thereof as long as the antisense DRP-1 sequence is capable of hybridizing to "sense" DRP-1 mRNA to prevent its translation into DRP-1 protein. Accordingly, an "antisense" sequence of the present invention is defined as a sequence which is capable of specifically hybridizing to "sense" DRP-1 mRNA to form a non-translatable double-stranded RNA molecule.

The antisense DRP-1 sequence need not hybridize to the entire length of the DRP-1 mRNA. Instead, it may hybridize to selected regions, such as the 5'-untranslated sequence, the coding sequence, or the 3'-untranslated sequence of the "sense" mRNA. In view of the size of the mammalian genome, the antisense DRP-1 sequence is preferably at least 17, more preferably at least 30, base pairs in length. However, shorter sequences may still be useful, i.e., they either fortuitously do not hybridize to other mammalian sequences, or such "cross-hybridization" does not interfere with the metabolism of the cell in a manner and to a degree which prevents the accomplishment of an object of this invention. The greater the length of the antisense sequence and the greater the number of complementary base pairs, the greater the number of non-complementary bases that can be tolerated, especially if the non-complementary bases are scattered. Both the preferred hybridization target on DRP-1 and the preferred antisense sequence length are readily determined by systematic experiment.

Standard methods such as described in Sambrooke et al., (1989) can be used to systematically remove an increasingly larger portion of the antisense DRP-1 sequence from a plasmid vector. Besides the full length antisense DRP-1 sequence, a series of staggered deletions may be generated, preferably at the 5'-end of the antisense DRP-1 sequence. This creates a set of truncated antisense DRP-1 sequences that still remain complementary to

preferably the 5'-end of the sense DRP-1 mRNA and as a result, still forms a RNA molecule that is double-stranded at the 5'-end of the sense DRP-1 mRNA (complements the 3'-end of an antisense DRP-1 RNA) and remains non-translatable.

The antisense RNA according to the present invention can be used in a method
5 to neutralize a mRNA molecule, which is the transcription product of the DNA sequence encoding the DRP-1 protein of SEQ ID NO:2, by allowing the antisense RNA to hybridize to the DRP-1 mRNA to prevent its translation into DRP-1 protein.

A further aspect of the present invention is directed to a composition, such as a pharmaceutical composition, which contains DRP-1, functional homologues or fragments
10 thereof and a pharmaceutically-acceptable excipient, carrier, diluent, or auxiliary agent.

An antibody, which specifically recognizes DRP-1 or functional homologues thereof is part of the present invention as long as the antibody does not cross-react with DAP-Kinase or ZIP-kinase. For instance, an antibody that specifically recognizes the unique 40 amino acid C-terminal tail of DRP-1, which is not present in DAP-Kinase or ZIP-kinase, is a
15 preferred embodiment of the antibody according to the present invention. Such an antibody can be used for diagnostic imaging, purification of DRP-1 etc.

The terms "antibody" or "antibodies" as used herein are intended to include intact antibodies, such as polyclonal antibodies or monoclonal antibodies (mAbs), as well as proteolytic fragments thereof such as the Fab or F(ab')₂ fragments. Furthermore, the DNA
20 encoding the variable region of the antibody can be inserted into other antibodies to produce chimeric antibodies (see, for example, U.S. Patent 4,816,567) or into T-cell receptors to produce T-cells with the same broad specificity (Eshhar et al., 1990; Gross, et al., 1989). Single chain antibodies can also be produced and used. Single chain antibodies can be single chain composite polypeptides having antigen binding capabilities and comprising a pair of
25 amino acid sequences homologous or analogous to the variable regions of an immunoglobulin light and heavy chain (linked V_H-V_L or single chain F_V). Both V_H and V_L may copy natural monoclonal antibody sequences or one or both of the chains may comprise a CDR-FR construct of the type described in U.S. Patent 5,091,513 (the entire contents of which are hereby incorporated herein by reference). The separate polypeptides analogous to the variable
30 regions of the light and heavy chains are held together by a polypeptide linker. Methods of production of such single chain antibodies, particularly where the DNA encoding the polypeptide structures of the V_H and V_L chains are known, may be accomplished in

accordance with the methods described, for example, in U.S. Patents 4,946,778, 5,091,513 and 5,096,815, the entire contents of each of which are hereby incorporated herein by reference.

As mentioned above, the terms "antibody" or "antibodies" are also meant to include both intact molecules as well as fragments thereof, such as, for example, Fab and $F(ab')_2$, which are capable of binding antigen. Fab and $F(ab')_2$ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl et al., 1983). It will be appreciated that Fab and $F(ab')_2$ and other fragments of the antibodies useful in the present invention may be used for the detection and quantitation of DRP-1 or functional homologues thereof according to the methods used for intact antibody molecules. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce $F(ab')_2$ fragments).

The present invention comprehends not only the intact antibodies or fragments, but also any molecule which includes an antigen binding portion of an antibody such that the molecule is capable of binding to the antigen. It is well within the skill of the art for the artisan to make e.g., fusion proteins which include antigen binding portions of an antibody fused to any other material which is desired to be carried to the antigen binding site, such as marker molecules, toxins, etc.

The antibodies, or fragments of antibodies, of the present invention may be used to quantitatively or qualitatively detect the presence of DRP-1 or functional homologues according to the present invention in a sample. The antibody according to the present invention may also be used for the isolation and purification of DRP-1 or homologues and fragments thereof, such as in an affinity column where the antibodies are immobilized on a solid phase support or carrier.

By "solid phase support or carrier" is intended any support capable of binding antigen or antibodies. Well-known supports, or carriers, include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a

bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

5 One of the ways in which the DRP-1-specific antibody can be detectably labeled is by linking the same to an enzyme and used in an enzyme immunoassay (EIA). This enzyme, in turn, when later exposed to an appropriate substrate, will react with the substrate in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means. The detection can be accomplished by
10 colorimetric methods which employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it
15 is possible to detect DRP-1 protein through the use of a radioimmunoassay (RIA) (Chard, T., "An Introduction to Radioimmune Assay and Related Techniques" (In: Work, T.S., *et al.*, Laboratory Techniques in Biochemistry in Molecular Biology, North Holland Publishing Company, New York (1978), incorporated by reference herein). The radioactive isotope can be detected by such means as the use of a gamma counter or a liquid scintillation counter or by
20 autoradiography. Radioactively labeled antibodies or antibody fragments can also be used for their capacity to kill cells bound by such antibodies, or cells in the immediate vicinity which are exposed to the radiation from such antibodies. It is also possible to label the antibody with a fluorescent compound, a chemiluminescent or bioluminescent compound.

The antibody molecules of the present invention may also be adapted for
25 utilization in an immunometric assay (also known as a "two-site" or "sandwich" assay) which is well known in the art.

In the present specification, the term "programmed cell death" is used to denote a physiological type of cell death which results from activation of some cellular mechanisms, i.e., death which is controlled by the cell's machinery. Programmed cell death
30 may, for example, be the result of activation of the cell machinery by an external trigger, e.g., a cytokine, which leads to cell death. The term "apoptosis" is also used interchangeably with programmed cell death.

The term "tumor" in the present specification denotes an uncontrolled growing mass of abnormal cells. This term includes both primary tumors, which may be benign or malignant, as well as secondary tumors, or metastases, which have spread to other sites in the body.

5 DRP-1 can be used to inhibit growth and metastasis of tumors. Tumor cells are exposed to a variety of death-inducing signals which, in combination with DAP-kinase-related I, can lead to death of the tumor cells. For example, in the blood stream, invading tumor cells must resist programmed cell death that is induced by interactions with cytotoxic T lymphocytes, natural killer cells, and macrophages, and with the cytokines which
10 these hematopoietic cells secrete (e.g., IFNs, TNF, IL-1 β). Tumor cells must also resist the apoptotic cell death induced by nitric oxide anions produced by the endothelial cells, and withstand mechanical shearing forces caused by hemodynamic turbulence. Moreover, during the intravasation or extravasation processes, and during growth in a foreign hostile microenvironment, locally produced inhibitory cytokines (e.g., TGF- β or loss of cell-matrix
15 interactions (e.g., detachment from the basement membranes) also trigger apoptotic cell death.

DRP-1 is useful in promoting death of tumor cells. The protein may be administered to patients, in particular, to cancer patients, which administration may cause death of the tumor cells. The protein may be administered *per se*, or may be administered by an expression vector comprising a DNA molecule of the present invention.

20 Because DRP-1 displays 80% identity with the catalytic domain of DAP-kinase and has a region which displays a high homology to the calmodulin-regulatory region of DAP-kinase, it is expected that DRP-1 has enzymatic kinase activity, which is calmodulin-dependent. Thus, DRP-1 has use as an enzyme and may be used, for example, as the enzyme in any *in vitro* enzymatic reaction which requires the presence of a kinase enzyme.
25 Accordingly, DRP-1 can be used *in vitro* to catalyze phosphorylation reactions as a kinase.

DRP-1 is capable of inducing apoptotic cell death when overexpressed in various cell lines. This ectopic cell death is blocked specifically by the death domain of DAP-kinase, suggesting possible crosstalk between these two kinases. Thus, DRP-1 may also be used for promoting the death of normal or tumor cells and for suppressing the metastatic
30 activity of tumor cells. A particular application of the death-promoting aspect is in therapy of diseases or disorders associated with uncontrolled, pathological cell growth, e.g., cancer (primary tumors and metastasis), psoriasis, autoimmune disease and others. Indeed, it is

expected that the DAP-kinase-related protein I of the present invention and DNA encoding it, may be used in the same manner as disclosed in detail in U.S. applications 08/810,712 and 08/631,097, as well as WO 95/10630.

According to a further aspect of the present invention, referred to herein at
5 times as "the screening aspect", DRP-1 DNA molecules are used in order to screen individuals for predisposition to cancer. In accordance with this aspect the screening is carried out by comparing the sequence of each of the DAP-kinase-related I DNA molecules to each of the respective DAP genes in the individual, or by following RNA and/or protein expression. The
10 absence of a DAP-kinase-related I gene, a partial deletion or any other difference in the sequence that indicates a mutation in an essential region, or the lack of a DRP-1 RNA and/or protein which may result in a loss of function may lead to a predisposition for cancer. For screening, preferably a battery of related DAP and DRP-1 genes maybe used, as well as different antibodies.

In the screening aspect, DAP-kinase related product I molecules may also be
15 used for prognostic purposes. For example, if a tumor cell lacks DRP-1 activity, this may reflect high chances of developing metastasis. In addition, DRP-1 positive cells may be more susceptible to control by chemotherapeutic drugs that work by inducing apoptosis, so that the choice of treatment modalities may be made based upon the DRP-1 state of the cells.

The DAP-kinase-related product can be used to screen individuals for
20 predisposition to cancer. There is provided a method for detecting the absence of a DRP-1 gene, a partial deletion or a mutation (i.e., point mutation, deletion or any other mutation) in the DRP-1 genes of an individual, or the absence of a DRP-1 RNA or protein, comprising probing genomic DNA, cDNA, or RNA from the individual with a DNA probe or a multitude of DNA probes having a complete or partial sequence of the DRP-1 genes, or probing protein
25 extracts with specific antibodies.

A particular application of the screening aspect of this invention is in the screening for individuals having a predisposition to cancer, an absence of the gene, or a detected mutation or deletion indicating that the individual has such a predisposition.

One example of a method in accordance with the screening aspect typically
30 comprises the following steps:

(a) obtaining a sample of either genomic DNA from cells of the individual or cDNA produced from mRNA of said cells;

(b) adding one or more DNA probes, each of said probes comprising a complete or partial sequence of a DRP-1 gene;

(c) providing conditions for hybridization to determine whether the DRP-1 gene is present or absent, i.e., whether there is a match between the sequence of the DNA probe or probes and a sequence in the DNA of said sample or a mismatch, a mismatch indicating a deletion or a mutation in the endogenous DNA and a predisposition to cancer in the tested individual.

Other examples of the screening aspect of the invention are well known to the skilled artisan and include, but are not limited to, Northern blots, RNase protection assays, and various PCR procedures.

The mutation in the DRP-1 gene, indicating a possible predisposition to cancer, can also be detected by the aid of appropriate antibodies which are able to distinguish between a mutated, a non-functional and a normal functional DRP-1 gene product. In addition, mutations that abolish protein translation or transcription due to promoter inactivation can be detected with the aid of antibodies that are used to react with protein cell extracts. Screening is also possible with respect to metastases.

Having now generally described the invention, the same will be more readily understood through reference to the following example which is provided by way of illustration and is not intended to be limiting of the present invention.

EXAMPLE

In this study, the identification and the structure/function analysis of a novel DAP-kinase-related protein, DRP-1, is described, DRP-1 is a 42kDa Ca^{2+} /CaM-regulated serine/threonine kinase which shows high degree of homology to DAP (Death Associated Protein)-kinase. The homology spans over the catalytic domain and the calmodulin-regulatory region, whereas the rest C-terminal part of the protein differs completely from DAP-kinase and displays no homology to any known protein. The catalytic domain is also homologous to the recently identified ZIP-kinase and to a lesser extent to the catalytic domains of DRAK1/2, thus forming together a novel subfamily of serine/threonine kinases. DRP-1 is localized to the cytoplasm as shown by immunostaining and cellular fractionation assays. *In vitro* kinase assays indicate that wild type DRP-1, but not a kinase inactive mutant, undergoes autophosphorylation and phosphorylates an external substrate in a Ca^{2+} /CaM-dependent

anner. Ectopically expressed DRP-1 is able to induce apoptosis in various types of cells. All killing by DRP-1 is dependent on two features: the intact kinase activity and the presence of C-terminal 40 amino acids shown to be involved in self-dimerization of the kinase. Interestingly, further deletion of the CaM-regulatory region overrode the indispensable role of C-terminal tail and generated a "super-killer" mutant. Finally, a dominant negative mutant of DAP-kinase encompassing the death domain is a potent blocker of apoptosis induced by DRP-1. This implies a possible functional connection between DAP-kinase and DRP-1. The experiments conducted in this study and the results obtained are presented

RESULTS AND METHODS

Cloning and Northern Blot Analysis

A PCR fragment of 364 bp was obtained from a λ gt11 human spleen cDNA library (Clontech) using primers from the deduced DRP-1 sequence, 5'-ATGCCGGATGAGGACCTGAGG-1066 (SEQ ID NO:13) and 3'-TCCACACTCCCACCCCAGACTC-1390 (SEQ ID NO:14). To obtain the full length cDNA of DRP-1, the same cDNA library was screened with the radiolabeled PCR product. Positive phage clones were isolated, cDNA was subcloned into a BlueScript vector and analyzed by restriction enzyme mapping and DNA sequencing. A 270 bp 3'-fragment from the full length cDNA of DRP-1 was generated by EcoRI digestion, and used to probe polyA⁺ RNA prepared by a standard procedure from various cell lines.

In vitro Transcription and Translation Assay

The full length cDNA was used as a template for *in vitro* transcription from the T7 promoter. This RNA was translated in reticulocyte lysate (TNT® T7 Quick Coupled Transcription/Translation System; Promega) by conventional procedures, with [³⁵S] methionine (Amersham) as a labeled precursor. The reaction product was then run on 12% SDS-PAGE gel, followed by sodium salicylate incubation for signal amplification. The gel was dried and exposed to X-ray film.

In vitro Kinase Assay

293 cells were transfected by a FLAG-tagged wild type DRP-1, DRP-1 K42A mutant, or mock-transfected. Cell lysates of 293 transfected cells were prepared as previously described (Deiss et al., 1995). Immunoprecipitation of DRP-1 or DRP-1 K42A mutant from 150µg total extract was done with 20µl anti-FLAG M2 gel (IBI, Kodak) in 500µl of PLB supplemented with protease and phosphatase inhibitors for 2h at 4°C. Following three washes with PLB, the immunoprecipitates were washed once with reaction buffer (50 mM HEPES pH 7.5, 20 mM MgCl₂, and 0.1 mg/ml BSA). The proteins bound to the beads were incubated for 15 min at 30°C in 50 µl of reaction buffer containing 15 µCi [γ 32p] ATP (3 pmole), 50µM ATP, 5µg MLC (Sigma), and where indicated, also 1µM bovine calmodulin (Sigma), 0.5 mM CaCl₂, or 3mM EGTA in the absence of calmodulin/CaCl₂. Protein sample buffer was added to terminate the reaction, and after boiling, the proteins were analyzed on 11% SDS-PAGE. The gel was blotted onto a nitrocellulose membrane and ³²P- labeled proteins were visualized by autoradiography.

15 Immunostaining of Cells

DRP-1 transfected or mock-transfected COS-7 cells were plated on glass cover-slips (13 mm diam.). After 48 hours, the cells were fixed/permeabilized in 3% formaldehyde for 5 min, methanol 5 min, acetone 2 min. The cells were blocked in 10% NGS for 30 min and incubated with anti-FLAG antibodies (dilution 1:100; IBI, Kodak) in 10% NGS for 60 min. Rhodamine-conjugated goat anti-mouse secondary antibodies (dilution 1:200; Jackson Immuno Research Lab.) and the nucleic acid dye, Oligreen (dilution 1:5000; Molecular Probes), for nuclear staining were then applied. The coverslips were mounted in Mowiol and observed under fluorescence microscope.

25 Detergent Extraction Assay

Sub-confluent cultures of COS-7 transfected cells, grown on 9 cm plate, were washed once with PBS and then with MES buffer (50 mM MES pH 6.8, 2.5 mM EGTA, 2.5 mM MgCl₂). The cells were extracted for 3 min with 0.5 ml of 0.5% Triton X-100 in MES buffer supplemented with protease inhibitors. The supernatant (the soluble fraction- Sol) was collected, centrifuged for 2 min. at 16,000x g at 4°C, and the clear supernatant was then transferred to new tubes. Two volumes of cold ethanol were added and the tubes were incubated at -20°C for overnight, centrifuged 10 min. at 16,000x g at 4°C and resuspended in

200 μ l of 2x protein sample buffer without dye. The detergent insoluble matrix (InSol) remaining on the plate was extracted in 200 μ l of 2x protein sample buffer, scraped from the plate with rubber policeman and collected into tube. The samples were loaded on 10% SDS-PAGE, 100 μ g protein extracts were loaded on each lane from the Sol fraction,

- 5 equivalent volumes of InSol were loaded. Analysis of the proteins was done using monoclonal anti-FLAG antibodies (dilution 1:200; IBI, Kodak).

Cell Lines, Transfections and Apoptotic Assays

- All cell lines were grown in DMEM (Biological Industries) supplemented with 10% fetal calf serum (Bio-Lab). For transient transfection, 1x10⁵ cells per well, were seeded in a 6 well plate a day before transfection. Transfections were done by calcium-phosphate method. For cell death assays by inducing overexpression, a mixture containing 1.5 μ g of cell death plasmid (expressing either DRP-1 or Δ CaM DAPk mutant) and 0.5 μ g of pEGFP-NI plasmid (Clontech) was used. For cell death protection assays we used a mixture containing 15 1.2 μ g of cell death inducing plasmid (either DRP-1 or Δ CaM DAPk mutant), 0.5 μ g of a plasmid to be tested for cell death protection (expressing DAPk-DD, DN FADD or luciferase as negative control), and 0.5 μ g of pEGFP-NI plasmid. Cells were counted and photographed 24 hours after transfection. In each transfection, three fields, each consisting of at least 100 GFP-positive cells, were scored for apoptotic cells according to their morphology. When indicated, cell lysates were prepared from the transient transfection at 24 hours, for protein analysis. The transfections of Rat embryo fibroblasts (REF) and FACS analysis of transfected fibroblasts for DNA content distribution were done as previously described (Kissil et al., 20 1998).

25 Co-immunoprecipitation assays

- 293 cells grown in 90mm plates (1x10⁶ cells/plate) were co-transfected with 5 μ g FLAG-tagged or HA-tagged DRP-1 and 20 μ g of HA-tagged or FLAG-tagged RFX1- Δ SmaI, respectively, or with DRP-1-HA and DRP-1-FLAG, 5 μ g each. Immunoprecipitation of DRP-1 or RFX1- Δ SmaI from 1mg total extract was done using anti-FLAG M2 gel or anti-30 HA as described above. Detection of bound proteins was done using anti-HA antibodies (dilution 1:1000, Babco) or anti-FLAG antibodies. For the deletion mutant study, 5 μ g of FLAG-tagged fully length DRP-1 were co-transfected with 5 μ g of HA-tagged DRP-1 deletion

mutants. Immunoprecipitation of DRP-1 from 1mg total extract was done using anti-FLAG M2 gel as described above. Detection of co-immunoprecipitated proteins (the mutants of DRP-1 or full length DRP-1) was done using anti-HA antibodies.

5 Nucleotide sequence accession number

The nucleotide sequence of human DRP-1 has been submitted to the GenBank™/EBI Data Bank (accession no. AF052941). The murine DRP-1 is also deposited at the GenBank™/EBI Data Bank (accession no. AF052942).

10 RESULTS

Cloning of DRP-1

To identify proteins that share homologous sequences with DAP-kinase, EST databases were searched using the BLAST™ program. Two ESTs of human and murine origin showed remarkable amino acid homology to the catalytic domains of DAP-kinase and the recently identified protein ZIP-kinase (79.5% and 80.2% identity, respectively). A second EST search was performed using the 5' and the 3' ends of the human EST, which identified a few more overlapping ESTs. A putative novel cDNA sequence was generated and used to design primers for cloning the full length cDNA. PCR performed on human spleen cDNA library amplified a 364 bp fragment that was further used to screen the same library. The full length cDNA was then isolated, subcloned into BlueScript vector, and sequenced.

The isolated cDNA was found to be 1742 bp long and to contain a serine/threonine kinase domain with all of the 12 characterized subdomains present (Park et al., 1997, Fig. 1A). Sequence alignment indicated that the catalytic domain of DRP-1 has 80% sequence identity to that of DAP-kinase and ZIP-kinase, yet less 50% sequence identity to the newly identified DRAK proteins (Fig. 2A). Like DAP-kinase but unlike ZIP-kinase, DRP-1 carries a typical CaM-regulatory region adjacent to its catalytic domain, as shown in Figures 1 and 2B. As compared with other kinases such as CaKIIa and MLCK, DRP-1 has the highest homology to DAP-kinase in this region, as shown in Figure 2B. The remaining short stretch of amino acids at the C-terminal part of DRP-1 (40 amino acid tail) displays no homology to any known protein.

Expression of DRP-1 and Tissue Distribution

To check the RNA expression of DRP-1, polyA+RNA was prepared from various cell lines and hybridized to a probe designed from the less conserved region of DRP-1. A single weak band of 1.9 kb appeared in some cell lines, in a Northern blot analysis of poly A+RNA (3 micrograms) extracted from various cell lines (Fig. 3A), suggesting that the mRNA is expressed at low amounts in HeLa, 293 and MCF-7 cells. The mRNA was hybridized with a radiolabeled human DRP-1 probe. The position of the transcript is indicated by an arrow. From PCR analysis of various cDNA libraries and the data gathered from EST searches, it was concluded that human DRP-1 is expressed, at least, in spleen, colon, breast, and leukocyte tissue.

In vitro transcription and translation assays conducted in reticulocyte lysates using the cloned DRP-1 cDNA as a template generated a single protein band of about 42 kDa in size, as predicted by its sequence. This protein band, obtained by Western blot analysis of *in vitro* transcribed and then translated DRP-1, is shown in Figure 3B. A FLAG-tagged DRP-1 was then cloned into pCDNA3 vector and expressed in HeLa cells. A protein of 42 kDa was evident upon immunoblot analysis of the cell lysates with anti-FLAG antibodies, shown in Figure 3C. In this case, 24 hours following transfection, the cells were harvested and lysed. The extracted proteins were separated by SDS-PAGE and then immunoblotted with anti-FLAG antibodies.

Cellular Localization of Ectopically Expressed DRP-1

In order to follow the cellular localization of the exogenous DRP-1, FLAG-tagged DRP-1 was expressed in COS-7 cells. COS-7 cells were transfected by a FLAG-tagged DRP-1 cloned in pCDNA3 vector, fixed and permeabilized in 1% formaldehyde followed by methanol/acetone treatment. Cells were visualized under fluorescence microscope. Immunoblot analysis proved that DRP-1 was expressed in these cells. For the immunostaining procedure, the non-transfected (Fig. 4A) and DRP-1 transfected (Fig. 4B) COS-7 cells were then fixed and reacted both with Oligreen for nuclear staining and anti-FLAG antibodies for DRP-1 staining. Specific DRP-1 staining was detected in the cytoplasm of these cells, as shown in Figure 4B.

A gentle cell extraction was performed with nonionic detergent, 0.5% TRITON X-100, that removes lipids and soluble proteins, leaving intact the detergent insoluble matrix composed of the nucleus, the cytoskeleton framework, and cytoskeleton-

associated proteins. In contrast to DAP-kinase, which is exclusively localized to the cytoskeleton, and hence found only in detergent insoluble fractions (Cohen et al., 1997, Fig. 4C), DRP-1 was preferentially eluted from the detergent soluble fraction, while a small amount was eluted from the insoluble fraction, as shown in Figure 4C. Thus, it was concluded that DRP-1 is a cytoplasmic protein with minor association with insoluble matrix components.

Intrinsic Kinase Activity of DRP-1

To test whether DRP-1 functions as a kinase as predicted from the amino acid sequence, an *in vitro* kinase assay was performed using myosin light chain (MLC) as an exogenous substrate. This substrate was chosen because it is phosphorylated by DAP-kinase (Cohen et al., 1997). DRP-1 was transfected into human kidney 293 cells, immunoprecipitated, and incubated with MLC in the presence and absence of Ca^{2+} and calmodulin. Both MLC phosphorylation and DRP-1 autophosphorylation were evident, as can be seen from Figure 5A.

In assaying the *in vitro* kinase activity of DRP-1, the proteins were assayed in the presence or absence of Ca^{2+} /CaM and MLC. The proteins were run on 11% SDS-PAGE and blotted to nitrocellulose membrane. Figure 5A shows the autophosphorylation of DRP-1 and MLC phosphorylation, respectively, as seen after exposure of X-ray film. Figure 5B shows the DRP-1 proteins by incubation of the same blot with anti-FLAG antibodies and ECL detection.

The addition of Ca^{2+} /calmodulin to the reaction mixture increased the amount of phosphorylated MLC, in accordance with the assumption that, like DAP-kinase, DRP-1 is negatively regulated by the autoinhibitory calmodulin binding domain, and that this inhibition is removed by the binding of Ca^{2+} /calmodulin. A catalytically inactive mutant of DRP-1, DRP-1 K42A, did not phosphorylate MLC and failed to undergo autophosphorylation even though higher amounts of DRP-1 protein were present, as can be seen from Figure 5A. Thus, DRP-1 was found to function *in vitro* as a kinase that is capable of phosphorylating itself and an external substrate. This latter property is stimulated by the addition of Ca^{2+} and calmodulin.

DRP-1 Induces Apoptosis in a Variety of Cell Lines

The high homology to DAP-kinase in the kinase and calmodulin-binding regions suggested the value of checking whether DRP-1 is involved in apoptosis. The wild type DRP-1 and the catalytically inactive mutant of DRP-1, DRP-1 K42A, which are cloned in pCDNA3 vector, were transfected into 293 cells. To quantitate the number of apoptotic cells, these constructs were transfected with a vector expressing the GFP protein. The GFP protein was used as a marker to visualize the transfected cells and to assess the apoptotic frequency among the transfectants according to morphological alterations. Apoptotic cells were scored after 24 hours. Overexpression of the DRP-1 resulted in massive apoptotic cell death (50-60%), as compared to the basal level of apoptotic cells caused by transfection of the non-relevant gene luciferase, shown in Figures 6A-6B and 7.

Most of the GFP positive green cells rounded up and shrunk; some of them showed cytoplasmic blebs, and some were further fragmented into "apoptotic bodies." In addition, some of the transfected cells detached from the plate. This apoptotic cell death was only slightly lower than that of an activated DAP- kinase mutant lacking the autoinhibitory calmodulin regulatory region (Δ CaM; apoptotic values of 70-80%). In contrast, when the cells were transfected with the kinase inactive mutant of DRP-1, DRP-1 K42A, as shown in Figures 6A-6D and 7, no apoptosis was observed. This experiment was repeated six times with reproducible results.

Western blot analysis of transfected cells, using anti- FLAG antibodies, confirmed the expression of both the exogenous wild type and K42A mutant of DRP-1 (Figures 8A and 8B). Similar results were also observed in human SV-80 fibroblasts. In another type of assay, the effect of ectopically expressed DRP-1 on the DNA content of rat embryo primary fibroblasts (REF cells) was assessed, as previously described (Kissil et al., 1999). The REFS were co-transfected with DRP-1 and a membrane-bound form of GFP and then after 48 hours subjected to FACS analysis of their DNA content. A fraction of cells displaying a sub-G1 population, indicative of cells containing fragmented DNA, appeared exclusively in the DRP-1 transfected cells but not in cells transfected with a control vector or with the DRP-1 K42A mutant form. No effect was found on cell cycle distribution of the viable cells.

To obtain the results shown in Figures 6A-6D, 1×10^5 293 cells/well were co-transfected with FLAG-tagged wild type DRP-1 or K42A mutant of DRP-1, 1.5 microgram/well and GFP, 0.5 microgram/well. GFP positive cells were visualized under

fluorescent microscope and scored for the appearance of apoptotic morphology 24 hours after transfection. Apoptotic cells are indicated by arrows. The fluorescent microscopic images correspond to 293 cells transfected by pCDNA3-luciferase as negative control (Fig. 6A), pCDNA3- Δ CaM DAP-kinase as positive control (Fig. 6B), pCDNA3-DRP-1 (Fig. 6C), pCDNA3-K42A DRP-1 (Fig. 6D).

In Figure 7, graphs show the percentage of apoptotic cells resulting from the above-mentioned transfections (average \pm S.D. calculated from triplicates of 100 cells each). The scores were taken from the same experiment shown in Figures 6A-6D.

In Figures 8A and 8B, proteins extracted from the transfected cells were run on 10% SDS-PAGE and blotted to nitrocellulose membrane. The blot was probed with anti-FLAG antibodies for DRP-1 detection and anti-vinculin antibodies to estimate the loaded protein amounts. The proteins were prepared from the same experiments shown in Figures 6A-6D.

Kinase Death Domain Protects From DRP-1 Induced Apoptosis

The structural homology of DRP-1 to DAP-kinase, the common regulation by Ca^{2+} /calmodulin, and the finding that both proteins caused apoptosis upon overexpression, suggested that they function along a common apoptotic pathway. In order to test this possibility, the effect of the dominant-negative DAP-kinase death domain (DAPk DD) on DRP-1-induced cell death was analyzed. The laboratory of the present inventor showed recently that overexpression of the fragment encompassing the death domain of DAP-kinase acts as a specific dominant-negative mutant, negating the effects of the full length protein (Datta et al., 1997). As a consequence, it protected cells from TNF- α , Fas and FADD/MORT1-induced cell death (Datta et al., 1997).

It has now been discovered that DAPk DD protected cell death induced by DRP-1 in 293 cells. As shown in Figure 9A, the apoptotic ratio dropped from 64.3% to 24.7%. A control transfection including DRP-1 and a non-relevant luciferase DNA excluded the possibility that this blockage was simply due to larger amount of DNA used in the transfection. Moreover, the effect of DAPk DD was specific, since the death domain of FADD failed to manifest a similar effect. (Figure 9A). Western blot analysis of transfected cells using anti-FLAG antibodies confirmed the expression of the exogenous DRP-1 in all transfections, as shown in Figure 9B. This experiment was repeated three times with

reproducible results. The ability of the death domain of DAP-kinase to block death induced by DRP-1 implies that DAP-kinase and DRP-1 function along a common pathway.

To obtain the results shown in Figure 9A, 1×10^5 cells/well of 293 cells were co-transfected with 1.2 microgram/well of FLAG-tagged wildtype DRP-1 and 0.5

5 microgram/well of GFP. The scores are the percentage of apoptotic cells given as average \pm S.D. and calculated from triplicates of 100 cells each.

To demonstrate the DRP-1 protein expression in 293 transfected cells shown in Figure 9B, proteins extracted from the transfected cells were separated on 10% SDS-PAGE and blotted to nitrocellulose membrane. The blot was hybridized with anti-FLAG antibodies
10 for DRP-1 detection and anti-vinculin antibodies to quantitate the loaded protein amounts. The proteins were prepared from the same experiment shown in Figure 9A.

Deletion of the C-terminal tail of DRP-1 abolishes its apoptotic activity, while further truncation of the CaM-regulatory region strongly enhances the apoptotic effect

15 In order to further understand the mode of DRP-1 action in apoptosis, constructs containing C-terminal truncations of DRP-1 tagged by HA were constructed (Fig. 10A). DRP-1 $\Delta 40$ lacks the most C-terminal part of DRP-1 which displays no homology to any known protein. DRP-1 $\Delta 73$ lacks, in addition to that, the CaM-regulatory region of DRP-1, and DRP-1 $\Delta 85$ contains only the catalytic domain. The wild type DRP-1 and the various
20 truncation mutants of DRP-1 were transfected into 293 cells. Induction of apoptotic cell death was assayed as mentioned above in DRP-1 induced apoptosis. Overexpression of the wild type DRP-1 resulted in apoptosis (25%) while the DRP-1 $\Delta 40$ had no effect in these assays. On the other hand, further truncations of the CaM-regulatory region, yielded mutants ($\Delta 73$, $\Delta 85$) which acted as "super-killers" (~90% apoptosis) (Figs. 11A and 11B). This
25 experiment was repeated three times with reproducible results. Western blot analysis of transfected cells, using anti-HA antibodies confirmed the expression of all DRP-1 forms (Fig. 10B). Thus, the dependence of the apoptotic effect of DRP-1 on its kinase activity was confirmed again, since removal of the CaM-regulatory region which acts as an autoinhibitory domain generates a constitutively active kinase. In addition, the existence of a positive
30 module in the C-terminal region of DRP-1, which is necessary for its pro-apoptotic effect, provided that the CaM-regulatory effect is still present, is shown.

The C-terminal part of DRP-1 functions as a homo-dimerization domain

Western analysis performed on proteins extracted from 293 cells transfected by FLAG-tagged DRP-1 revealed an additional band (not shown). This observation led the present inventor to test whether DRP-1 can undergo homo-dimerization. To this aim, two constructs expressing DRP-1 fused to either FLAG or HA tags were co-transfected into 293 cells and classical pull-down experiments with each one of the two epitopes were performed. FLAG-tagged DRP-1 could be shown to bind specifically to HA-tagged DRP-1 in both IP directions (Fig. 12A, see lane 3 in both IP Panels). No binding of DRP-1-HA to FLAG beads or to the irrelevant cytoplasmic protein RFX- Δ SmaI could be observed (Fig. 12A, see IP anti-FLAG panel, lanes 2 or 1+2, respectively). Also non-specific binding of DRP-1-FLAG to HA bead or to RFX- Δ SmaI protein could not be detected (Fig. 12A, see IP anti-HA panel, lanes 1 or 1+2, respectively). Western analyses confirmed the expression of all proteins in these cell extracts (Fig. 12A, see Western panels).

The observation that a C-terminal truncation of 40 amino acids in DRP-1 abolished its apoptotic effect upon ectopic expression in 293 cells, prompted the present inventor to test whether this domain may be involved in the homo-dimerization of DRP-1. DRP-1-FLAG was co-expressed in conjugation with the various deletion mutants of DRP-1 tagged by HA. A strong binding of DRP-1-FLAG to the wild type DRP-1-HA was detected, whereas the binding to DRP-1 Δ 40 was mostly abolished (Fig. 12B, upper IP panel, compare lane 1 to 2-4). Western analysis confirmed the expression of wild type DRP-1-HA and all other DRP-1-HA deletion mutants in these transfections (Fig. 12B, see Western panel). Lower IP panel confirmed the expression of wild type DRP-1-FLAG in all these transfections. Thus, the present inventor concluded that a region spanning the C-terminal 40 amino acids of DRP-1 is responsible for its homo-dimerization. This homo-dimerization is probably required for the apoptotic effect of DRP-1, since DRP-1- Δ 40 has lost the ability to induce apoptosis in 293 cells (Figs. 11A and 11B).

Having now fully described this invention, it will be appreciated by those skilled in the art that the same can be performed within a wide range of equivalent parameters, concentrations, and conditions without departing from the spirit and scope of the invention and without undue experimentation.

While this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This

application is intended to cover any variations, uses or adaptations of the inventions following, in general, the principles of the invention and including such departures from the present disclosure as come within know or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth as follows in the scope of the appended claims.

All references cited herein, including journal articles or abstracts, published or unpublished U.S. or foreign patent application, issued U.S. or foreign patents, or any other references, are entirely incorporated by reference herein, including all data, tables, figures, and text presented in the cited references. Additionally, the entire contents of the references cited within the references cited herein are also entirely incorporated by reference.

Reference to known method steps, conventional method steps, known methods or conventional methods is not in any way an admission that any aspect, description or embodiment of the present invention is disclosed, taught or suggested in the relevant art.

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art (including the contents of the references cited herein), readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance presented herein, in combination with the knowledge of one of ordinary skill in the art.

REFERENCES

- Anderson, P., "Kinase cascades regulating entry into apoptosis," Microbiol. Mo. Biol. Rev., **61**, 33-46, 1997.
- Ausubel et al., Current Protocols in Molecular Biology, Green Publications and Wiley
5 Interscience, New York, 1987-1999.
- Basu, S., and Kolesnick, R., "Stress signals for apoptosis: ceramide and c-Jun kinase,"
Oncogene, **17**, 3277-3285, 1998.
- Bokoch, G.M., "Caspase-mediated activation of PAK2 during apoptosis: proteolytic kinase
activation as a general mechanism of apoptotic signal transduction?", Cell death diff.,
10 **5**, 637-645, 1998.
- Cardone, M.H., Salveson, G.S., Widmann, C., Johnson, G. And Frisch, S.M., "The
regulation of anoikis: MEKK-1 activation requires cleavage by caspases.", Cell,
90, 315-323, 1997.
- Cardone, M.H., Roy, N., Stennicke, H.R., Salvesen, G.S., Franke, T.F., Stanbridge, E.,
15 Frisch, S., and Reed, J.S., "Regulation of cell death protease caspase-9 by
phosphorylation.", Science, **282**, 318-321, 1998.
- Cohen, O., Feinstein, E., and Kimchi, A., "DAP-kinase is a Ca²⁺/calmodulin-dependent,
cytoskeletal-associated protein kinase, with cell death-inducing functions that depend
on its catalytic activity.", EMBO J., **16**, 998-1008, 1997.
- 20 Cohen, O., Inbal, B., Kissil, J.L., Feinstein, E., Spivak, T., and Kimchi, A., "DAP-kinase
participates in TNF- α and Fas-induced apoptosis and its function requires the death
domain.", J. Cell. Biol., in press, 1999.
- Datta, S.R., Dudek, H., Tao, X., Masters, S., Fu, H., Gotoh, Y., and Greenberg, M.E., "Akt
phosphorylation of BAD couples survival signals to the cell-intrinsic death
25 machinery.", Cell, **91**, 231-241, 1997.
- Deiss, L.P., Feinstein, E., Berissi, H., Cohen, O., and Kimchi, A., "Identification of a novel
serine/threonine kinase and a novel 15-kD protein as potential mediators of the
gamma interferon-induced cell death.", Genes Dev., **9**, 15-30, 1995.
- Deiss, L.P. and Kimchi, A., "A genetic tool used to identify thioredoxin as a mediator of a
30 growth inhibitory signal.", Science, **252**, 117-120, 1991.

- del Peso, L., Gonzalez-Garcia, M., Page, C., Herrera, R., and Nunez, G., "Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt.", Science, **282**, 318-321, 1997.
- Eshhar, Z. et al., Br. J. Cancer Suppl., **10**, 27-9, 1990.
- 5 Feinstein, E., Kimchi, A., Wallach, D., Boldin, M., and Varfolomeev, E., "The death domain: a module shared by proteins with diverse cellular functions.", Trends Biochem. Sci., **20**, 342-344, 1995.
- Green, D., and Kroemer, G., "The central executioners of apoptosis: caspases or mitochondria?" Trends Cell Biol., **8**, 267-271, 1998.
- 10 Gross, G. et al., Proc. Natl. Acad. Sci. USA, **86**, 10024-8, 1989.
- Hanks, S.K., and Quinn, A.M., "Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members.", Methods Enzymol., **200**, 38-62, 1991.
- Inbal, B., Cohen, O., Polak-Charcon, S., Kopolovic, J., Vadai, E., Eisenbach, L., and Kimchi, A., "DAP kinase links the control of apoptosis to metastasis.", Nature, **390**, 180-184, 1997.
- 15 Inbal, B., Kissil, J.K., Cohen, O., Spivak-Kroizman, T., and Kimchi, A., "The DAP-related protein kinases-a novel subfamily of serine/threonine kinases with a possible link to apoptosis.", submitted, 1999.
- 20 Jacobson, M.D., Weil, M., and Raff, M.C., "Programmed cell death in animal development." Cell, **88**, 347-354, 1997.
- Kawai, T., Matsumoto, M., Takeda, K., Sanjo, H., and Akira, S., "ZIP kinase, a novel serine/threonine kinase which mediates apoptosis.", Mol. Cell Biol., **18**, 1642-1651, 1998.
- Kelliher, M.A., Grimm, S., Ishida, Y., Kuo, F., Stanger, B.Z., and Leder, P., "The death domain kinase RIP mediates the TNF-induced NF-kappaB signal.", Immunity, **8**, 297-303, 1998.
- 25 Kimchi, A., J. Cell. Biochem., **50**, 1-9, 1992.
- Kimchi, A., "DAP genes: novel apoptotic genes isolated by a functional approach to gene cloning.", Biochim. Biophys. Acta, **1377**, F13-33, 1998.
- 30 Kissil, J.L., and Kimchi, A., "Death-associated proteins: from gene identification to the analysis of their apoptotic and tumor suppressive functions.", Mol. Med. Today, **4**, 268-74, 1998.

- Kissil, J.L., Cohen, O., Raveh, T., and Kimchi, A., "DAP-kinase loss of expression in various carcinoma and B-cell lymphoma cell lines: possible implications for role as tumor suppressor gene.", EMBO J., **18**, 353-362, 1999.
- Kogel, D., Plottner, O., Landsberg, G., Christian, S., and Scheidtmann, K.H., "Cloning and
5 characterization of Dlk, a novel serine/threonine kinase that is tightly associated with chromatin and phosphorylates core histones.", Oncogene, **17**, 2645-2654, 1998.
- Levy et al., Mol. Cell. Biol., **13**, 7942-7952, 1993.
- Levy-Strumpf, N., and Kimchi, A., "Death associated proteins (DAPs): from gene
10 identification to the analysis of their apoptotic and tumor suppressive functions."
Oncogene, **17**, 3331-3340, 1998.
- Maundrell, K., Antonsson, B., Magnenat, E., Camps, M., Muda, M., Chabert, C., Gillieron, C., Boschert, U., Vial-Knecht, E., Martinou, J.C., and Artkinstall, S., "Bcl-2
undergoes phosphorylation by c-Jun N-terminal kinase/stress-activated protein
15 kinases in the presence of the constitutively active GTP-binding protein Rac1.", J. Biol. Chem., **272**, 25238-25342, 1997.
- McCarthy, J.V., Ni, J., and Dixit, V.M., "RIP2 is a novel NF-kappaB-activating and cell death-inducing kinase, J. Biol. Chem., **273**, 16968-75, 1998.
- Meinkoth et al., Anal. Biochem., **138**, 267-284, 1984.
- Park, J., Kim, I., Oh, Y.J., Lee, K., Han, P.L., and Choi, E.J., "Activation of c-Jun N-
20 terminal kinase antagonizes an anti-apoptotic action of Bcl-2.", J. Biol. Chem., **272**, 16725-16728, 1997.
- Peitenpol et al., Cell, **61**, 777-785, 1990.
- Sambrooke et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY, 1989.
- 25 Sanjo, H., Kawai, T., and Akira, S., "DRAKS, novel serine/threonine kinases related to death-associated protein kinase that trigger apoptosis.", J. Biol. Chem., **273**, 29066-29071, 1998.
- Stanger, B.Z., Leder, P., Lee, T.H., Kim, E., and Seed, B., "RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell
30 death.", Cell, **81**, 513-523, 1995.
- Sun, X., Lee, J., Navas, T., Baldwin, D.T., Stewart, T.A., and Dixit, V.M., "RIP3, a Novel Apoptosis-inducing Kinase.", J. Biol. Chem., **274**, 16871-16875, 1999.

- Tompkins, J.D., Higgins, D.G., Gibson, T., J., "CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice.", Nucleic Acids Res., **22**, 4673-4680, 1994.
- Verheij, M., Ruiter, G.A., Zerp, S.F., Van Blitterswijk, W.J., Fuks, Z. Haimovitz-Friedman, A., and Bartelink, H., "The role of the stress-activated protein kinase (SAPK/JNK) signaling pathway in radiation-induced apoptosis.", Radiother. Oncol., **47**, 225-232, 1998.
- et al., J. Nucl. Med., **24**, 316-325, 1983.
- et al., E., "Life, death and the pursuit of apoptosis.", Genes Dev., **10**, 1-15, 1996.
- et al., X., Khosravi-Far, R., Chang, H.Y., and Baltimore, D., Cell, **89**, 1067-1076, 1997.
- et al., W., Huang, B.C., Shen, M., Quast, J., Chan, E., Xu, X, Nolan, G.P., Payan, D.G. and Luo, Y., "Identification of RIP3, a RIP-like kinase that activates apoptosis and Nf-kappa B.", Curr. Biol., **9**, 539-42, 1999.

WHAT IS CLAIMED IS:

1. An isolated polypeptide, which is a calmodulin-dependent serine/threonine kinase, or a fragment thereof, selected from the group consisting of:

5 (A) a polypeptide which is capable of inducing cell death (apoptosis) and comprises the amino acid sequence of SEQ ID NO:2;

(B) a polypeptide which has a property being capable of inducing cell death and has at least 85% sequence identity to the amino acid sequence of SEQ ID NO:2;

(C) a fragment of a polypeptide of SEQ ID NO:2 which is capable of inducing cell death;

10 (D) a fragment which is capable of inducing cell death and has at least 85% sequence identity to fragment (C);

(E) a fragment of a polypeptide of SEQ ID NO:2 which lacks the property of being capable of inducing cell death and which inhibits the ability of polypeptide (A) or (B) to induce cell death; and

15 (F) a fragment which lacks the property of being capable of inducing cell death and which inhibits the ability of polypeptide (A) or (B) to induce cell death, said fragment having at least 85% sequence identity to fragment (E).

2. An isolated DNA molecule comprising a nucleotide sequence encoding the polypeptide or fragment thereof according to claim 1.

20 3. The isolated DNA molecule according to claim 1, wherein said nucleotide sequence encodes the amino acid sequence of SEQ ID NO:2.

4. The isolated DNA molecule according to claim 3, wherein said nucleotide sequence corresponds to nucleotides 62 to 1141 of SEQ ID NO:1.

5. The isolated DNA molecule according to claim 3, which consists of the nucleotide sequence corresponding to nucleotides 62 to 1141 of SEQ ID NO:1.

6. An isolated DNA molecule which hybridizes to the DNA molecule of claim 5 under moderately stringent conditions and encodes a calmodulin-dependent serine/threonine kinase having the property of being capable of inducing cell death.

7. An isolated DNA molecule which hybridizes to the DNA molecule of claim 5 under highly stringent conditions and encodes a calmodulin-dependent serine/threonine kinase having the property of being capable of inducing cell death.

8. A polypeptide consisting of an amino acid sequence selected from the group consisting of amino acid residues 13 to 275 of SEQ ID NO:2 and an amino acid sequence having at least 85% sequence identity to residues 13 to 275 of SEQ ID NO:2.

9. An isolated DNA molecule comprising a nucleotide sequence encoding the polypeptide of claim 8.

10. The isolated DNA molecule according to claim 9, wherein said nucleotide sequence encodes the amino acid sequence corresponding to residues 13 to 275 of SEQ ID NO:2.

11. The isolated DNA molecule according to claim 10, wherein said nucleotide sequence is selected from the group consisting of nucleotides 98 to 886 of SEQ ID NO:1 and a nucleotide sequence which hybridizes to nucleotides 98 to 886 of SEQ ID NO:1 under moderately stringent conditions.

12. The isolated DNA molecule according to claim 11, wherein said nucleotide sequence hybridizes to nucleotides 98 to 886 of SEQ ID NO:1 under highly stringent conditions.

13. A polypeptide consisting of an amino acid sequence selected from the group consisting of amino acid residues 321 to 360 of SEQ ID NO:2 and an amino acid sequence having at least 85% sequence identity to residues 321 to 360 of SEQ ID NO:2.

14. An isolated DNA molecule comprising a nucleotide sequence encoding the
5 polypeptide of claim 13.

15. The isolated DNA molecule according to claim 14, wherein said nucleotide sequence encodes the amino acid sequence corresponding to residues 321 to 360 of SEQ ID NO:2.

16. The isolated DNA molecule according to claim 15, wherein said nucleotide
10 sequence is selected from the group consisting of nucleotides 1022 to 1141 of SEQ ID NO:1 and a nucleotide sequence which hybridizes to nucleotides 1022 to 1141 of SEQ ID NO:1 under moderately stringent conditions.

17. The isolated DNA molecule according to claim 16, wherein said nucleotide
15 sequence hybridizes to nucleotides 1022 to 1141 of SEQ ID NO:1 under highly stringent conditions.

18. A vector comprising the isolated DNA molecule according to any of claims 2-7, 9-12 and 14-17.

19. A host cell transformed with the isolated DNA molecule according to any of claims 2-7, 9-12, and 14-17.

20. A composition comprising a polypeptide according to any one of claims 1, 8 and 13, and a pharmaceutically acceptable excipient, carrier, diluent or auxiliary agent.

21. A molecule containing an antigen binding portion of an antibody which specifically recognizes the polypeptide according to any one of claims 1, 8 and 13 with the proviso that said antibody does not cross-react with DAP kinase or ZIP kinase.

22. The antibody according to claim 21, which is a monoclonal antibody.

23. A single stranded RNA molecule complementary to at least a portion of the isolated messenger RNA molecule which is the transcription product of the DNA sequence encoding a polypeptide of SEQ ID NO:2, wherein said complementary single stranded RNA molecule is capable of hybridizing to said isolated messenger RNA to prevent its translation
30 into said polypeptide of SEQ ID NO:2.

24. A method of neutralizing a messenger RNA molecule, which is the transcription product of the DNA sequence encoding a polypeptide of SEQ ID NO:2, comprising the step

of contacting the single stranded RNA molecule of claim 23 with the messenger RNA to neutralize the messenger RNA by hybridizing thereto and preventing its translation into the polypeptide of SEQ ID NO:2.

25. A method for screening individuals for a predisposition to cancer comprising the steps of:

(a) obtaining a sample of either genomic DNA from cells of the individual or cDNA produced from mRNA of said cells; and

(b) determining if there is a mutation in the nucleotide sequence of the DRP-1 gene.

26. The method according to claim 25 wherein a mutation in the nucleotide sequence of DRP-1 is determined by a process comprising the steps of:

(a) adding one or more nucleic acid probes to the sample of genomic DNA or cDNA, wherein each probe comprises a portion of the nucleotide sequence of DRP-1;

(b) providing conditions for hybridization between the nucleic acid probe or probes and the DNA of said samples; and

(c) determining on the basis of the hybridization whether there is a match between the sequence of the nucleic acid probe or probes and a sequence in the DNA of said sample, or whether there is a mismatch, a deletion or a mutation in the genomic DNA or cDNA and a predisposition to cancer of the tested individual.

```

GACCGCGGCAGCTCAGCCTCCCGCGATTGTATGTTCCAGCCCTCAATGAGGAGTCCAAA 60
M E P F K Q Q K V E D F Y D I G E E L G 20
CATTGAGCCATTCAAGCAGCAGAGGTGGAGGACTTTTATGACATCGGAGAGGAGCTGGG 120
S G Q F A I V K K C R E K S T G L E Y A 40
GAGTGGCCAGTTTGGCATCTGTGAAGAAGTCCCGGAGAGAGCAGCGGGCTTGAATATGC 180
A K F I K K R Q S R A S R R G V S R E E 60
AGCCAAAGTTTCATCAAGAAAGCGGAGAGCGCGGAGCCCGCGGGTGTGAGCGGGAGGA 240
I E R E V S I L R Q V L H H N V I T L H 80
GATCGAGCGGGAGGTGAGCATCTCGCGCAAGTGTCTGACCCACAATGTTCATCAGCTGCA 300
D V Y E N R T D V V L I L E L V S G G E 100
CGACGTCTATGAGAACCGGACCGGAGCTGTGTCTCATCTTGAAGTAGTGTCTGGAGGAGA 360
L F D F L A Q K E S L S E E E A T S F I 120
GCTCTTGGATTTCCTGGCCAGAAAGGTCTACTGAGTGAAGAGGAGGCGGACAGCTTCAT 420
K Q I L D G V N Y L R T K K I A H F D L 140
TAAGCAGATCTCGATGGGTGAACTACCTTCACACAAAGAAAATTGCTCACTTTGATCT 480
K P E N I M L L D K N I P I P H I K L I 160
CAAGCCAGAAAACATTATGTTGTAGACAGAAATATCCCATTCACACATCAAGCTGAT 540
D F G L A H E I E D G V E F K N I P G T 180
TGACTTTGGTCTGGCTCAGAAATAGAAATGGAGTGAATTTAAGAAATATTTTGGGAC 600
P E F V A P E I V N Y E P L G L E A D M 200
GCGGAAATTTGTGCTCCAGAAATTTGAACTACGAGCGCCCTGGGTCTGGAGGCTGACAT 660
W S I G V I T Y I L L S G A S P F L G D 220
GTGGAGCATAGGCGTCATCAGCTACATCTCTTAAGTGGAGCATCCCTTTCTGGGAGA 720
T K Q E T L A N I T S V S Y D F D E E F 240
CAGCAAGCAGGAACACTGGCAAAATATCAGATCAGTGAATTTAGGACTTTGATGAGGAATT 780
F S H T S E L A K D F I R K L L V K E T 260
CTTCAGCCATACGAGCGAGCTGGCCAAAGGACTTTATTCGGAAGCTTCTGTTTAAAGAGAC 840
R K R L T I Q E A L R H P W I T P V D N 280
CGGAAACGGCTCACAATCCAGAGGCTCTCAGACACCCCTGGATCACCGCGGTGACAA 900
Q O A M V R R E S V V N L E N F R K Q Y 300
CCAGCAAGCCATGTTGCGCAGGAGTTTGTGCTCAATTTGGAGAACTTACGAAAGGACTA 960
V R R R W K L S F S I V S L C N H L T R 320
TGTCGGCAGGCTGTGGAGCTTTCTTTCAGCAATCTGCTCTCTTTCAGGAGCTTACCCG 1020
S L M K K V H L R P D E D L R N C E S D 340
CTCGCTGATGAGAAGGTGCACCTGAGGCGGATGAGGACCTGAGGAACGTGAGAGTGA 1080
T E E D I A R R K A L H P R R R S S T S 360
CACTGAGGAGGACATGCCAGGAGAAAGCCCTCCACCCAGGAGGAGGAGCAGCAGCTC 1140
CTATCTGGCCTGACCTGCACTGGCCGCGAGGAGTTTGGGCCAGCGGGCTCCCTTCT 1200
GTGCAGACTTTTGGACCCAGCTCAGCAGCAGCAGCCGGCGCTCTGAGCACTTTGCAAGA 1260
GAGATGGGCCCAAGGAATTCAGAAAGCTTCCAGGCAAGCCAGGAGACCTGGGAGCTGT 1320
GGCTGTCTCTGTGGAGGAGGCTCCAGCATTTCCCAAAGCTCTTAATTCTCCATAAAATGG 1380
GCTTTCTCTGTCTGCCATCTCAGAGTCTGGGTGGGAGTGTGGAATTAGGAAAACAAT 1440
ATAAAGGACATCTCATCATCACGGGTGAAGGTGAGAGTAAAGCAGCTTCTTCACAGG 1500
CTGAGGGGGTTTCAGAACAGCTGGCCAAAATTTACACCAGAGAGACAGAGTCTCCCA 1560
TTGGGAACAGGGTGATTGAGGAAAGTGAACTTGGGTGTGAGGGACCAATCTGTGACCT 1620
CCCAGAACCATGGAAGCCAGGAGCTCAGGCTGACCAACACCTCAGACCTTCTGAAGCAGC 1680
CCATTGTGGCCCGCATGTTGTAATTTTCTCATTTTATTAAAGcttctggtttacctg 1740
aa 1742

```

FIG. 1

DAP-kinase	13	YDTGEELGSGGF	AVVKKCHERKST	GLQYAAKFIK
ZIP-kinase	13	YEMGEELGSGGF	AVVKKCHERKST	GLQYAAKFIK
DRP-1	13	YDIQEELGSGGF	AVVKKCHERKST	GLQYAAKFIK
DRAK1	61	LCPSRELGSGGF	AVVKKCHERKST	GLQYAAKFIK
DRAK2	33	ILTSKELGSGGF	AVVKKCHERKST	GLQYAAKFIK
DAP-kinase	46	KRRFKS	SRRGVSRE	DIHERVSTLKEN - OHPNVT
ZIP-kinase	46	KRRLLSS	RRGVSRE	DIHERVSTLKEN - RHPNVI
DRP-1	46	KRRSH	RRGVSRE	DIHERVSTLKEN - RHPNVI
DRAK1	94	KRR	RRGVSRE	DIHERVSTLKEN - RHPNVI
DRAK2	66	KRR	RRGVSRE	DIHERVSTLKEN - RHPNVI
DAP-kinase	78	TLHEVYEN	ETAS	EMVEEYAAQGEI
ZIP-kinase	78	TLHEVYEN	ETAS	EMVEEYAAQGEI
DRP-1	78	TLHEVYEN	ETAS	EMVEEYAAQGEI
DRAK1	123	TLHEVYEN	ETAS	EMVEEYAAQGEI
DRAK2	95	TLHEVYEN	ETAS	EMVEEYAAQGEI
DAP-kinase	109	ESLSE	EPATSH	IKRGLDGVN
ZIP-kinase	109	ESLSE	EPATSH	IKRGLDGVN
DRP-1	109	ESLSE	EPATSH	IKRGLDGVN
DRAK1	156	EMVSE	ENDVIR	LKCRNE
DRAK2	128	EMVSE	ENDVIR	LKCRNE
DAP-kinase	142	PENIM	LDKN	MRNFRNKLI
ZIP-kinase	142	PENIM	LDKN	MRNFRNKLI
DRP-1	142	PENIM	LDKN	MRNFRNKLI
DRAK1	189	PONIC	ETSE	PLGD - EKV
DRAK2	161	PONIC	ETSE	PLGD - EKV
DAP-kinase	175	KNIF	GLPEE	VAPEN
ZIP-kinase	175	KNIF	GLPEE	VAPEN
DRP-1	175	KNIF	GLPEE	VAPEN
DRAK1	221	REIM	GLPEE	VAPEN
DRAK2	193	REIM	GLPEE	VAPEN
DAP-kinase	208	YIL	SSGAS	FRGDR
ZIP-kinase	208	YIL	SSGAS	FRGDR
DRP-1	208	YIL	SSGAS	FRGDR
DRAK1	254	YVME	TSIS	PPGND
DRAK2	226	YVME	TSIS	PPGND
DAP-kinase	241	FSNT	SA	DAKDE
ZIP-kinase	241	FSNT	SA	DAKDE
DRP-1	241	FSNT	SA	DAKDE
DRAK1	287	FDVL	SE	SAVD
DRAK2	259	FDVL	SE	SAVD
DAP-kinase	274	WL		
ZIP-kinase	274	WL		
DRP-1	274	WL		
DRAK1	320	WL		
DRAK2	292	WL		

FIG. 2A

DAP-kinase	1	NMEKEKK	FAARKWK	QSVRL	ESTCORLSR	29
DRP-1	1	NREN	FAARK	QSVRL	ESTCORLSR	29
smMLCK	1	SKDR	FAARK	QSVRL	ESTCORLSR	29
CaMKIIa	1	TVDC	FAARK	QSVRL	ESTCORLSR	29
CaMKI	1	VSE	FAARK	QSVRL	ESTCORLSR	29
CaMKIV	1	MDTAQ	FAARK	QSVRL	ESTCORLSR	29
ZIP-kinase	1	GEDSG	FAARK	QSVRL	ESTCORLSR	29

FIG. 2B




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 5A




FIG. 5B

FIG. 6B

FIG. 6D




FIG. 6A

FIG. 6C




FIG. 7




FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B




FIG. 10A






FIG. 10B

**FIG. 11A****FIG. 11B**




FIG. 12A




FIG. 12B

SEQUENCE LISTING

<110> KIMCHI, Adi
 MCINNIS A., Patricia
 YEDA RESEARCH AND DEVELOPMENT COMPANY LTD.

<120> DAP-KINASE RELATED PROTEIN

<130> KIMCHI2A

<140> 00

<141> 1999-06-15

<150> 60/089,294

<151> 1998-06-15

<160> 14

<170> PatentIn Ver. 2.0

<210> 1

<211> 1742

<212> DNA

<213> Human

<220>

<221> CDS

<222> (62)..(1141)

<400> 1

gaccgcggca gctcagcctc ccgccgattg tatgttccag gcctcaatga ggagtccaaa 60

c atg gag cca ttc aag cag cag aag gtg gag gac ttt tat gac atc gga 109

Met Glu Pro Phe Lys Gln Gln Lys Val Glu Asp Phe Tyr Asp Ile Gly

1 5 10 15

gag gag ctg ggg agt ggc cag ttt gcc atc gtg aag aag tgc cgg gag 157

Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Lys Lys Cys Arg Glu

20 25 30

aag agc acg ggg ctt gag tat gca gcc aag ttc atc aag aag cgg cag 205

Lys Ser Thr Gly Leu Glu Tyr Ala Ala Lys Phe Ile Lys Lys Arg Gln

35 40 45

agc cgg gcg agc cgg cgc ggt gtg agc cgg gag gag atc gag cgg gag 253

Ser Arg Ala Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu Arg Glu

50 55 60

2

gtt aaa gag acc cgg aaa cgg ctc aca atc caa gag gct ctc aga cac 877
 Val Lys Glu Thr Arg Lys Arg Leu Thr Ile Gln Glu Ala Leu Arg His
 260 265 270

ccc tgg atc acg cgg gtg gac aac cag caa gcc atg gtg cga cgg gag 925
 Pro Trp Ile Thr Pro Val Asp Asn Gln Gln Ala Met Val Arg Arg Glu
 275 280 285

tct gtg gtc aat ctg gag aac ttc agg aag cag tat gtc cgc agg cgg 973
 Ser Val Val Asn Leu Glu Asn Phe Arg Lys Gln Tyr Val Arg Arg Arg
 290 295 300

tgg aag ctt tcc ttc agc atc gtg tcc ctg tgc aac cac ctc acc cgc 1021
 Trp Lys Leu Ser Phe Ser Ile Val Ser Leu Cys Asn His Leu Thr Arg
 305 310 315 320

tcg ctg atg aag aag gtg cac ctg agg cgg gat gag gac ctg agg aac 1069
 Ser Leu Met Lys Lys Val His Leu Arg Pro Asp Glu Asp Leu Arg Asn
 325 330 335

tgt gag agt gac act gag gag gac atc gcc agg agg aaa gcc ctc cac 1117
 Cys Glu Ser Asp Thr Glu Glu Asp Ile Ala Arg Arg Lys Ala Leu His
 340 345 350

cca cgg agg agg agc agc acc tcc taactggcct gacctgcagt ggccgccagg 1171
 Pro Arg Arg Arg Ser Ser Thr Ser
 355 360

gaggtttggg cccagcgggg ctcccttctg tgcagacttt tggaccacag tcagcaccag 1231

caccggggcg tcttgagcac ttgcaagag agatggggccc aaggaattca gaagagcttg 1291

caggcaagcc aggagaccct gggagctgtg gctgtcttct gtggaggagg ctccagcatt 1351

cccaaagctc ttaattctcc ataaaatggg ctttcctctg tctgccatcc tcagagtctg 1411

ggggtgggagt gtggacttag gaaaacaata taaaggacat cctcatcatc acgggggtgaa 1471

ggtcagagta aggcagcctt cttcacaggc tgaggggggtt cagaaccagc ctggccaaaa 1531

attacaccag agagacagag tcctcccat tgggaacagg gtgattgagg aaagtgaacc 1591

ttgggtgtga gggaccaatc ctgtgacctc ccagaacat ggaagccagg acgtcaggct 1651

gaccaacacc tcagaccttc tgaagcagcc cattgctggc ccgccatgtt gtaattttgc 1711

tcatttttat taaacttctg gtttacctga a 1742

<210> 2

<211> 360

<212> PRT

<213> Human

<400> 2

Met Glu Pro Phe Lys Gln Gln Lys Val Glu Asp Phe Tyr Asp Ile Gly
 1 5 10 15

Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Lys Lys Cys Arg Glu
 20 25 30

Lys Ser Thr Gly Leu Glu Tyr Ala Ala Lys Phe Ile Lys Lys Arg Gln
 35 40 45

Ser Arg Ala Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu Arg Glu
 50 55 60

Val Ser Ile Leu Arg Gln Val Leu His His Asn Val Ile Thr Leu His
 65 70 75 80

Asp Val Tyr Glu Asn Arg Thr Asp Val Val His Ile Leu Glu Leu Val
 85 90 95

Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Gln Lys Glu Ser Leu Ser
 100 105 110

Glu Glu Glu Ala Thr Ser Phe Ile Lys Gln Ile Leu Asp Gly Val Asn
 115 120 125

Tyr Leu His Thr Lys Lys Ile Ala His Phe Asp Leu Lys Pro Glu Asn
 130 135 140

Ile Met Leu Leu Asp Lys Asn Ile Pro Ile Pro His Ile Lys Leu Ile
 145 150 155 160

Asp Phe Gly Leu Ala His Glu Ile Glu Asp Gly Val Glu Phe Lys Asn
 165 170 175

Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn Tyr Glu
 180 185 190

Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly Val Ile Thr Tyr
 195 200 205

Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Asp Thr Lys Gln Glu
 210 215 220

Thr Leu Ala Asn Ile Thr Ser Val Ser Tyr Asp Phe Asp Glu Glu Phe
 225 230 235 240

Phe Ser His Thr Ser Glu Leu Ala Lys Asp Phe Ile Arg Lys Leu Leu
 245 250 255

Val Lys Glu Thr Arg Lys Arg Leu Thr Ile Gln Glu Ala Leu Arg His
 260 265 270

Pro Trp Ile Thr Pro Val Asp Asn Gln Gln Ala Met Val Arg Arg Glu
 275 280 285

Ser Val Val Asn Leu Glu Asn Phe Arg Lys Gln Tyr Val Arg Arg Arg
 290 295 300

Trp Lys Leu Ser Phe Ser Ile Val Ser Leu Cys Asn His Leu Thr Arg
 305 310 315 320

Ser Leu Met Lys Lys Val His Leu Arg Pro Asp Glu Asp Leu Arg Asn
 325 330 335

Cys Glu Ser Asp Thr Glu Glu Asp Ile Ala Arg Arg Lys Ala Leu His
 340 345 350

Pro Arg Arg Arg Ser Ser Thr Ser
 355 360

<210> 3

<211> 263

<212> PRT

<213> Human

<400> 3

Tyr Asp Thr Gly Glu Glu Leu Gly Ser Gly Gln Phe Ala Val Val Lys
 1 5 10 15

Lys Cys Arg Glu Lys Ser Thr Gly Leu Gln Tyr Ala Ala Lys Phe Ile
 20 25 30

Lys Lys Arg Arg Thr Lys Ser Ser Arg Arg Gly Val Ser Arg Glu Asp
 35 40 45

Ile Glu Arg Glu Val Ser Ile Leu Lys Glu Ile Gln His Pro Asn Val
 50 55 60

Ile Thr Leu His Glu Val Tyr Glu Asn Lys Thr Asp Val Ile Leu Ile

<400> 4
Tyr Glu Met Gly Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Arg
1 5 10 15

Lys Cys Arg Gln Lys Gly Thr Gly Lys Glu Tyr Ala Ala Lys Phe Ile
 20 25 30
 Lys Lys Arg Arg Leu Ser Ser Ser Arg Arg Gly Val Ser Arg Glu Glu
 35 40 45
 Ile Glu Arg Glu Val Asn Ile Leu Arg Glu Ile Arg His Pro Asn Ile
 50 55 60
 Thr Leu His Asp Ile Phe Glu Asn Lys Thr Asp Val Val Leu Ile
 65 70 75 80
 Leu Glu Leu Val Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Glu Lys
 85 90 95
 Ser Leu Thr Glu Asp Glu Ala Thr Gln Phe Leu Lys Gln Ile Leu
 100 105 110
 Gly Val His Tyr Leu His Ser Lys Arg Ile Ala His Phe Asp Leu
 115 120 125
 Pro Glu Asn Ile Met Leu Leu Asn Lys Asn Val Pro Asn Pro Arg
 130 135 140
 Ile Lys Leu Ile Asp Phe Gly Ile Ala His Lys Ile Glu Ala Gly Asn
 145 150 155 160
 Glu Phe Lys Asn Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile
 165 170 175
 Val Asn Tyr Glu Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly
 180 185 190
 Val Ile Thr Tyr Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Glu
 195 200 205
 Thr Lys Gln Glu Thr Leu Thr Asn Ile Ser Ala Val Asn Tyr Asp Phe
 210 215 220
 Asp Glu Glu Tyr Phe Ser Asn Thr Ser Glu Leu Ala Lys Asp Phe Ile
 225 230 235 240
 Arg Arg Leu Leu Val Lys Asp Pro Lys Arg Arg Met Thr Ile Ala Gln
 245 250 255
 Ser Leu Glu His Ser Trp Ile
 260

<210> 5

<211> 261

<212> PRT

<213> Human

<400> 5

Leu Cys Pro Gly Arg Glu Leu Gly Arg Gly Lys Phe Ala Val Val Arg
 1 5 10 15

Lys Cys Ile Lys Lys Asp Ser Gly Lys Glu Phe Ala Ala Lys Phe Met
 20 25 30

Arg Lys Arg Arg Lys Gly Gln Asp Cys Arg Met Glu Ile Ile His Glu
 35 40 45

Ile Ala Val Leu Glu Leu Ala Gln Asp Asn Pro Trp Val Ile Asn Leu
 50 55 60

His Glu Val Tyr Glu Thr Ala Ser Glu Met Ile Leu Val Leu Glu Tyr
 65 70 75 80

Ala Ala Gly Gly Glu Ile Phe Asp Gln Cys Val Ala Asp Arg Glu Glu
 85 90 95

Ala Phe Lys Glu Lys Asp Val Gln Arg Leu Met Arg Gln Ile Leu Glu
 100 105 110

Gly Val His Phe Leu His Thr Arg Asp Val Val His Leu Asp Leu Lys
 115 120 125

Pro Gln Asn Ile Leu Leu Thr Ser Glu Ser Pro Leu Gly Asp Ile Lys
 130 135 140

Ile Val Asp Phe Gly Leu Ser Arg Ile Leu Lys Asn Ser Glu Glu Leu
 145 150 155 160

Arg Glu Ile Met Gly Thr Pro Glu Tyr Val Ala Pro Glu Ile Leu Ser
 165 170 175

Tyr Asp Pro Ile Ser Met Ala Thr Asp Met Trp Ser Ile Gly Val Leu
 180 185 190

Thr Tyr Val Met Leu Thr Gly Ile Ser Pro Phe Leu Gly Asn Asp Lys
 195 200 205

Gln Glu Thr Phe Leu Asn Ile Ser Gln Met Asn Leu Ser Tyr Ser Glu
 210 215 220

Glu Glu Phe Asp Val Leu Ser Glu Ser Ala Val Asp Phe Ile Arg Thr
 225 230 235 240

Leu Leu Val Lys Lys Pro Glu Asp Arg Ala Thr Ala Glu Glu Cys Leu
 245 250 255

Lys His Pro Trp Leu
 260

<210> 6

<211> 261

<212> PRT

<213> Human

<400> 6

Ile Leu Thr Ser Lys Glu Leu Gly Arg Gly Lys Phe Ala Val Val Arg
 1 5 10 15

Gln Cys Ile Ser Lys Ser Thr Gly Gln Glu Tyr Ala Ala Lys Phe Leu
 20 25 30

Lys Lys Arg Arg Arg Gly Gln Asp Cys Arg Ala Glu Ile Leu His Glu
 35 40 45

Ile Ala Val Leu Glu Leu Ala Lys Ser Cys Pro Arg Val Ile Asn Leu
 50 55 60

His Glu Val Tyr Glu Asn Thr Ser Glu Ile Ile Leu Ile Leu Glu Tyr
 65 70 75 80

Ala Ala Gly Gly Glu Ile Phe Ser Leu Cys Leu Pro Glu Leu Ala Glu
 85 90 95

Met Val Ser Glu Asn Asp Val Ile Arg Leu Ile Lys Gln Ile Leu Glu
 100 105 110

Gly Val Tyr Tyr Leu His Gln Asn Asn Ile Val His Leu Asp Leu Lys
 115 120 125

Pro Gln Asn Ile Leu Leu Ser Ser Ile Tyr Pro Leu Gly Asp Ile Lys
 130 135 140

Ile Val Asp Phe Gly Met Ser Arg Lys Ile Gly His Ala Cys Glu Leu
 145 150 155 160

Arg Glu Ile Met Gly Thr Pro Glu Tyr Leu Ala Pro Glu Ile Leu Asn

[illegible]

```
<210> 7
<211> 29
<212> PRT
<213> Human
```

```
<400> 7
Asn Met Glu Lys Phe Lys Lys Phe Ala Ala Arg Lys Lys Trp Lys Gln
  1             5             10            15
```

Ser Val Arg Leu Ile Ser Leu Cys Gln Arg Leu Ser Arg
20 25

```
<210> 8
<211> 29
<212> PRT
<213> Human
```

<400> 8
Asn Leu Glu Asn Phe Arg Lys Gln Tyr Val Arg Arg Arg Trp Lys Leu
1 5 10 15

Ser Phe Ser Ile Val Ser Leu Cys Asn His Leu Thr Arg
20 25

<210> 9

<211> 29
<212> PRT
<213> Human

<400> 9
Thr Cys Asp Cys Leu Lys Lys Leu Asn Ala Arg Arg Lys Leu Lys Gly
1 5 10 15

Ala Ile Leu Thr Thr Met Leu Ala Thr Arg Asn Phe Ser
20 25

<210> 10
<211> 28
<212> PRT
<213> Human

<400> 10
Val Ser Glu Gln Ile Lys Lys Asn Phe Ala Lys Ser Lys Trp Lys Gln
1 5 10 15

Ala Phe Asn Ala Thr Ala Val Val Arg His Met Arg
20 25

<210> 11
<211> 32
<212> PRT
<213> Human

<400> 11
Met Asp Thr Ala Gln Lys Lys Leu Gln Glu Phe Asn Ala Arg Arg Lys
1 5 10 15

Leu Lys Ala Ala Val Lys Ala Val Val Ala Ser Ser Arg Leu Gly Ser
20 25 30

<210> 12
<211> 28
<212> PRT
<213> Human

<400> 12
Gly Glu Asp Ser Gly Arg Lys Pro Glu Arg Arg Arg Leu Lys Thr Thr

1

5

10

15

Arg Leu Lys Glu Tyr Thr Ile Lys Ser His Ser Ser
20 25

<210> 13

<211> 20

<212> DNA

<213> Human

<400> 13

ggccggatga ggacctgagg

20

<210> 14

<211> 21

<212> DNA

<213> Human

<400> 14

tccacatccc accccagact c

21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 9/12, 1/20, 5/00, 15/00, C12Q 1/68; C07H 21/04; A61K 38/51

US CL : 435/194, 320.2, 325, 252.3, 6; 424/94.5; 536/23.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/194, 320.2, 325, 252.3, 6; 424/94.5; 536/23.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9510630 A (YEDA RESEARCH AND DEVELOPMENT CO. LTD.) 20 April 1995 (20.04.95), see the entire article specially compare their kinase domain (AA 13-275) to a fragment of DRP-1 and its homologs in claims 1C-D.	1-7 and 20
X	DEISS et al. Identification of a novel serine/threonine kinase and a novel 15-kD protein as potential mediators of the γ interferon-induced cell death. Genes & Development. 01 January 1995, Vol. 9, pages 15-30, see figures 4 and 7.	8-12, 20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 NOVEMBER 1999

Date of mailing of the international search report

19 NOV 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MARYAM MONSHIPOURI CV

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

(continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Priority*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	<p>Database EST on GenCore version 4.5, Accession No. H27327, HILLIER et al. 'The WashU-Merck Project', definition: y116d12.r1 Soars breast 2NbHBst Homosapiens cDNA clone. 12 July 1995, see their residues 81-259 for 100% identity to residues 1300-1478 of SEQ ID NO:1.</p>	<p>23 ---- 24</p>

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-20, 23-24

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Detailed Reasons For Holding Lack of Unity of Invention:

This International Preliminary Examination Authority has found 3 inventions claimed in the International Application covered by the claims indicated below:

Group I, claims 1-20 and 23-24, drawn to isolated DNA molecules encoding calmodulin-dependent serine/threonine kinases (DRP-1) and their fragments, vectors and host cells comprising said DNA molecules, single stranded antisense RNA molecules complementary to at least a portion of the transcription product of said DNA molecules, methods of neutralizing a messenger RNA molecule, which is the transcription product of DRP-1 encoding gene comprising contacting said antisense RNA molecules to messenger RNA molecules in order to prevent translation, calmodulin-dependent serine/threonine (DRP-1) kinases and their fragments together with compositions comprising said kinases and their fragments.

Group II, claims 21-22, drawn to a molecule containing an antigen binding portion of an antibody which specifically binds said DRP-1 kinases.

Group III, claims 25-26, drawn to a method of screening cancer patients comprising obtaining a sample of either genomic DNA or cDNA of cancer cells and determining if a mutation occurred in the gene encoding DRP-1.

The inventions listed as Groups I-III do not relate to a single inventive concept because they are considered to be three categories of invention and are not drawn to combination of categories (i.e. categories 1-5), specified in 37 CFR section 1.475(B).



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/12, 1/20, 5/00, 15/00, C12Q 1/68, C07H 21/04, A61K 38/51	A1	(11) International Publication Number: WO 99/66030 (43) International Publication Date: 23 December 1999 (23.12.99)
(21) International Application Number: PCT/US99/13411 (22) International Filing Date: 15 June 1999 (15.06.99) (30) Priority Data: 60/089,294 15 June 1998 (15.06.98) US (71) Applicant (for all designated States except US): YEDA RESEARCH AND DEVELOPMENT COMPANY LTD. [IL/IL]; P.O. Box 96, 76100 Rehovot (IL). (71) Applicant (for SD only): MCINNIS, Patricia, A. [US/US]; Apartment #203, 2325 42nd Street N.W., Washington, DC 20007 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): KIMCHI, Adi [IL/IL]; 38 Hashalom Street, 43561 Raanana (IL). (74) Agent: BROWDY, Roger, L.; Browdy and Neimark, P.L.L.C., Suite 300, 419 Seventh Street N.W., Washington, DC 20004 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: DAP-KINASE RELATED PROTEIN (57) Abstract <p>A new protein, which is a novel homologue of DAP-kinase, has been isolated. This novel calmodulin-dependent kinase is a cell death-promoting protein functioning in the biochemical pathway which involves DAP (death-associated protein)-kinase (e.g., forming a cascade of sequential kinases, one directly activating the other). Alternatively, the two kinases may operate to promote cell death in parallel pathways.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DAP-KINASE RELATED PROTEIN**CROSS-REFERENCE TO RELATED APPLICATIONS**

5 The present application is related to and hereby incorporates by reference the entire contents of each of application serial no. 08/810,712, filed March 3, 1997, and application serial no. 08/631,097, filed April 12, 1996, the latter of which has been published as PCT publication WO 95/10630 on April 20, 1995. The present application also claims priority from provisional application 60/089,294, filed June 15, 1998, the entire contents of which are hereby
10 incorporated by reference.

BACKGROUND OF THE INVENTIONField of the Invention

 The present invention is directed to a DAP-kinase related protein.

Description of the Related Art

 One of the factors which determines the proliferation state of cells is the balance between the growth-promoting effects of proto-oncogenes and the growth-constraining effects of tumor-suppressor genes. One mechanism by which
15 tumor-suppressor genes exert their growth-constraining effect is by inducing a cell to undergo a physiological type of death. Such a controlled cell death is evident in a multitude of physiological conditions including metamorphosis, synaptogenesis of neurons, death of lymphocytes during receptor repertoire selection, and controlled homeostasis in the bone marrow and other proliferative tissues, etc. This cell death is regulated by the interaction of
20 the cell with other cells or with cell products, for example through the activity of suitable cytokines.

 Growth-inhibiting cytokines have a double effect on the target cell. They can either inhibit the proliferation of the cell and/or give rise to cell death. To date, blockage or activation of expression of known tumor-suppressor genes was shown to counteract or
25 enhance, respectively, cytokines inhibition of cells growth (Kimchi, 1992) but did not have any effect on the death- promoting action of cytokines. For example, the growth inhibitory response to cytokines, such as TGF- β , was markedly reduced by the inactivation of the Rb gene, or the response to IL-6 was enhanced by introducing activated p53 genes (Pietenpol et al., 1990; Levy et al, 1993).

Apoptosis is a genetically controlled cell death process which is important in various developmental stages, as well as for cell maintenance and tissue homeostasis (Jacobson et al., 1997). During the last few years, many of the key players in this process have been identified, including receptors, adaptor proteins, proteases, and other positive and negative regulators (Green et al., 1998; White, 1996). One of the positive mediators of apoptosis, which has been cloned by the present inventors, is DAP-kinase (Deiss et al., 1995). This protein was discovered by a functional approach to gene cloning, based on transfections of mammalian cells with anti-sense cDNA libraries and subsequent isolation of death-protective cDNA fragments (Deiss et al., 1995; Deiss et al., 1991; Kimchi, 1998; Kissil et al., 1998; Levy-Strumpf et al., 1998). The anti-sense cDNA of DAP-kinase protected HeLa cells from interferon-gamma-induced cell death, and this property served as the basis for its selection.

DAP-kinase is a calcium/calmodulin-regulated 160 kDa serine/threonine protein kinase associated with actin microfilaments (Deiss et al., 1995; Cohen et al., 1997). Its structure contains at least two additional domains that might mediate interactions with other proteins: ankyrin repeats, and a typical death domain located at the C-terminal part of the protein (Deiss et al., 1995; Cohen et al., 1997). Overexpression of DAP-kinase in various cell lines results in cell death, and this death-promoting effect of DAP-kinase depends on at least three features: the catalytic activity, presence of the death domain, and the correct intracellular localization (Cohen et al., 1997; Cohen et al., 1999). Several independent lines of evidence proved that DAP-kinase is involved in apoptosis triggered by different external signals, including interferon- γ , TNF- α , activated Fas receptors, and detachment of cells from the extracellular matrix (Deiss et al., 1995; Cohen et al., 1997; Cohen et al., 1999; Inbal et al., 1997). A tumor suppressive function was recently attributed to the DAP-Kinase, coupling the control of apoptosis to metastasis (Inbal et al., 1997).

So far, only a few serine/threonine kinases were implicated in the regulation of programmed cell death, either as death-promoting and death-protecting proteins (Anderson, 1997; Bokoch, 1998). One such candidate is the JNK/SAPK (Basu et al., 1998). In one example, it was shown to mediate apoptosis induced by detachment from extracellular matrix (named anoikis) (Cardone et al., 1997). In this system, the JNK pathway is activated by MEKK-1, whose kinase activity is stimulated by caspase cleavage (Cardone et al., 1997). JNK

may antagonize BCL-2 anti-apoptotic effects by phosphorylation (Park et al., 1997; Maundrell et al., 1997).

Another serine/threonine kinase is RIP, which like DAP-Kinase also possesses the death domain. RIP was shown to positively mediate apoptosis in cell cultures (Stanger et al., 1995). However, *in vivo* studies in RIP-deficient mice demonstrated its ability to exert anti-apoptotic effects by mediating the TNF- α -induced TNF- β activation (Kelliher et al., 1998). Other RIP members, RIP2 and RIP 3 were also recently identified and shown to possess pro-apoptotic effects (McCarthy et al., 1998; Sun et al., 1998; Yu et al., 1999).

Among the negative regulators of apoptosis is the protein kinase AKT. This protein was shown to phosphorylate BAD and thereby to prevent it from complexing and blocking the anti-apoptotic activity of BCL-X_L (Datta et al, 1997; del Peso et al., 1997). AKT was also recently shown to phosphorylate pro-caspase-9, thus blocking its normal processing (Cardone et al., 1998).

Recently, the isolation and characterization of novel kinase members, homologous in their catalytic domains to DAP-kinase, was reported (Kawai et al., 1998; Kogel et al., 1998; Sanjo et al., 1998). One protein, named ZIP-kinase, was found to be 80% identical to DAP-kinase within the kinase domain, yet it lacks the CaM-regulatory domain and the other domains and motifs characteristic of DAP-kinase. Zip-kinase contains a leucine zipper domain at the C-terminus and is localized to the nucleus (Kawai et al., 1998; Kogel et al. 1998). The activation of ZIP kinase occurs by a different mechanism involving homo-dimerization, mediated by its leucine zipper domain. However, unlike DAP-kinase, ZIP-kinase is a nuclear protein, which instead of being regulated by a calmodulin-binding domain, is activated by homo-dimerization of its leucine-zipper motifs (Kogel et al., 1998). Another two less conserved nuclear proteins, DRAK1 and DRAK2, which are closely related to each other, and which share 50% identity with the kinase domain of DAP-kinase, were also recently characterized. Like ZIP-kinase, the DRAK1 and DRAK2 proteins also lack the CaM-regulatory domain. The overexpression of these two proteins in NIH3T3 cells induces some morphological changes associated with apoptosis, dependent on the functionality of their kinase domain (Sanjo et al., 1998). Together these kinases form a novel subfamily of serine/threonine kinases, as is evident from multiple sequence and phylogenetic analysis (Inbal et al., 1999).

Ectopic expression of the three wild type kinases, but not their catalytically inactive mutants, induced morphological changes characteristic of apoptosis (Kawai et al., 1998; Sanjo et al., 1998). Yet, in the case of ZIP-Kinase, these results are still controversial (Kogel et al., 1998).

5 Citation of any document herein is not intended as an admission that such document is pertinent prior art, or considered material to the patentability of any claim of the present application. Any statement as to the content or a date of any document is based on information available to the applicant at the time of filing and does not constitute an admission as to the correctness of such a statement

SUMMARY OF THE INVENTION

10 A new protein, DAP-Kinase-related 1 protein (DRP-1), which is a novel homologue of DAP-kinase, has been isolated. This novel calmodulin-dependent kinase is a 42kDa serine/threonine kinase which shows a high degree of homology to DAP-kinase both in
15 its catalytic domain and its calmodulin-regulatory region. The catalytic domain of DRP-1 is also homologous to recently identified ZIP-kinase and, to a lesser extent, to the catalytic domains of DRAK1/2.

DRP-1 is localized to the cytoplasm as shown by immunostaining and cellular fractionation assays. *In vitro* kinase assays indicate that wild type DRP-1, but not a kinase
20 inactive mutant, undergoes autophosphorylation and phosphorylates an external substrate in a Ca^{2+} /CaM-dependent manner. Ectopically expressed DRP-1 is able to induce apoptosis in various types of cells; with this killing being dependent on its kinase activity. The dominant negative form of DAP-Kinase (DAPk DD) is a potent blocker of apoptosis induced DRP-1. Thus, DRP-1 may be a death-promoting protein functioning in the biochemical pathway which
25 involves DAP (death-associated protein)-kinase (e.g., forming a cascade of sequential kinases, one directly activating the other). Alternatively, the two kinases may operate to promote cell death in parallel pathways.

The present invention provides for a DRP-1 protein and functional homologues thereof having at least 85% sequence identity to the DRP-1 sequence of SEQ ID NO:2. Also
30 provided is a fragment of DRP-1, which either is capable of inducing cell death or lacks such capability but instead is capable of inhibiting the activity of DRP-1 or a functional homologue

thereof to induce cell death, and a homologous fragment which has at least 85% sequence identity thereto and which has the same properties.

The present invention further provides an isolated DNA molecule encoding for such DRP-1 protein, functional homologues thereof, or fragments thereof. Also included within the scope of the present invention are isolated DNA molecules which hybridize to the nucleotide sequence encoding DRP-1 protein under moderately or highly stringent conditions and encode a calmodulin-dependent serine/threonine kinase having the property of being capable of inducing cell death.

Other further aspects of the present invention include a composition comprising the DRP-1 protein, functional homologues and fragments thereof, and an antibody which specifically recognizes DRP-1 but does not cross-react with DAP kinase or ZIP kinase.

Yet another aspect of the present invention is directed to a single stranded RNA molecule complementary to at least a portion of the mRNA encoding the DRP-1 protein of SEQ ID NO:2. This single stranded antisense RNA molecule can be used in a method of neutralizing DRP-1 mRNA by hybridizing to the DRP-1 mRNA to prevent its translation into DRP-1 protein.

The present invention also provides a method for screening individuals for predisposition to cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide (SEQ ID NO:1) and amino acid (SEQ ID NO:2) sequence of the DAP-kinase homologue, DRP-1. The initiation (ATG) and stop (TAA) codons are boxed. The polyadenylation signal (ATTAAA) is underlined. The kinase domain and the calmodulin regulatory regions are in bold or underlined by a dash, respectively.

Figures 2A-2B show the multiple sequence alignment of the serine/threonine kinase domains (Fig. 2A) of the DAP-kinase-related proteins, DAP-kinase (SEQ ID NO:3), ZIP-kinase (SEQ ID NO:4), DRP-1 (corresponding to residues 13-275 of SEQ ID NO:2), DRAK1 (SEQ ID NO:5) and DRAK 2 (SEQ ID NO:6), conducted according to Hanks and Quinn (1991) with identical amino acids boxed and homologous amino acids shown with gray shading, and the multiple sequence alignment of the calmodulin regulatory regions (Fig. 2B) of DAP-kinase (SEQ ID NO:7), DRP-1 (corresponding to residues 292 to 320 of SEQ ID NO:2), smMLCK (SEQ ID NO:8), CaMKIIa (SEQ ID NO:9), CaMKI (SEQ ID NO:10),

CaMKIV (SEQ ID NO:11), and ZIP-Kinase (SEQ ID NO:12) conducted manually, keeping the conserved (boxed) regions aligned to each other. The corresponding region of ZIP-Kinase which does not contain homology to DAP-Kinase and ZIP-Kinase CAM-regulatory regions is given at the bottom of Fig. 2B.

Figure 3A shows Northern blot analysis of polyA+RNA extracted from various cell lines for mRNA expression of DRP-1, Figure 3B show Western blot analysis of *in vitro* transcription and translation of DRP-1, and Figure 3C shows protein expression of DRP-1 in HeLA cells on an immunoblot.

Figures 4A and 4B show control COS-7 cells and cellular localization of DRP-1 in COS-7 cells, respectively, and Figure 4C shows a Western blot of fractions from a detergent extraction of COS-7 cells transfected with a pCDNA3 vector expressing either FLAG-tagged DRP-1 or DAP-Kinase.

Figure 5A shows *in vitro* kinase activity of DRP-1 and Figure 5B shows a Western blot of DRP-1 proteins.

Figures 6A-6B show fluorescent microscope images of 293 cells transfected by pCDNA3-luciferase as a negative control (Fig. 6A), by pCDNA3- Δ CaM DAP-Kinase as positive control (Fig. 6B), by pCDNA -DRP-1 (Fig. 6C), and by pCDNA3-K42A DRP-1 (Fig. 6D). Apoptotic cells are indicated by arrows.

Figure 7 shows the scores of apoptotic cells in a graph of the percentage of apoptotic cells resulting from the transfections of Figs. 6A-6D.

Figures 8A and 8B show DRP-1 protein expression in 293 transfected cells in immunoblots to anti-FLAG antibodies (Fig. 8A) and anti-vinculin antibodies (Fig. 8B).

Figure 9A shows that DAP kinase death domain protects from DRP-1 induced apoptosis, and Figure 9B shows an immunoblot of DRP-1 protein expression in 293 transfected cells.

Figure 10A shows a schematic representation of a series of generated deletion mutant, and Figure 10B shows an immunoblot containing extracts of 293 cells transiently transfected with GFP and the series of deletion mutants, (DRP-1 fragments, cloned in pCDNA3, and tagged with HA epitope at the C-terminus), as in Figs. 8A and 8B are probed with anti-HA antibodies for DRP-1 detection and anti-vinculin antibodies to quantitate the loaded protein amounts. In Figure 10B, pCDNA3²-luciferase is the negative control.

Figure 11A shows fluorescent microscope images of the transiently transfected cells of Fig. 10B, and Figure 11B shows a graph of the score in percent apoptotic cells in Fig. 11A resulting from co-transfections of 293 cells with 1-2 μ g HA-tagged wild type DRP-1 or various deletion mutants of DRP-1 after 24 hours (average S.D. calculated from triplicates of 100 cells each).

Figures 12A and 12B show by Western analysis that the C-terminal part of DRP-1 is required for its homo-dimerization. In Figure 12A, wild type DRP-1 is shown to undergo specific homo-dimerization. The lanes correspond to the following co-transfections (5 μ g of DRP-1 constructs and 20 μ g of RFX1- Δ SmaI constructs/9mm plate): (1) DRP-1-FLAG+RFX1- Δ SmaI-HA (control to rule-out nonspecific attachment of DRP-1 to HA beads or to an irrelevant gene). (2) RFX- Δ SmaI-FLAG+DRP-1-HA (control to rule out nonspecific attachment of DRP-1 to FLAG beads or to an irrelevant gene). (3) DRP-1-FLAG+DRP-1-HA. Both IP directions and their Western blottings are shown. In Figure 12B, truncation of C-terminal 40 amino acids of DRP-1 is shown to abolish its homo-dimerization. The lanes correspond to the following co-transfections (5 μ g of each construct/90mm plate): (1) DRP-1-FLAG+DRP-1-HA (2) DRP-1-FLAG+DRP-1- Δ 40-HA (3) DRP-1-FLAG+DRP-1- Δ 73-HA (4) DRP-1-FLAG+DRP-1- Δ 85-HA. The lower panel quantitate the immunoprecipitation efficiency of DRP-1-FLAG by the anti-FLAG antibodies.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the discovery by the present inventor of a novel serine/threonine kinase with remarkable homology to the catalytic and CaM-regulatory domains of DAP-kinase. This kinase, named DAP-kinase-related protein 1 (DRP-1), is a 42kDa cytoplasmic protein which exhibits minor associations with insoluble matrix elements. The nucleotide (SEQ ID NO:1) and amino acid (SEQ ID NO:2) sequences of this DRP-1 protein are shown in Fig 1. It is composed of 1742 nucleotides. The predicted initiation and stop codons are boxed, and the polyadenylation signal is underlined. The protein kinase domain is shown in bold letters and corresponds to amino acid residues 13 to 275 of SEQ ID NO:2. This protein displays 80% identity with the catalytic domain of DAP-kinase. The calmodulin-regulatory region is underlined with a dashed line; this region displays high homology to the corresponding region in DAP-kinase. The remainder of the C-terminal short part of the protein (the last 40 amino acid residues corresponding to residues 321 to 360 of

SEQ ID NO:2) differs completely from DAP-kinase. Thus, DAP-kinase-related I protein does not carry all of the other motifs and protein modules characteristic of DAP-kinase. The mRNA expression levels transcribed from this gene are low.

Another protein, ZIP-kinase, which by virtue of its sequence homology to the kinase domain of DAP-Kinase, is also a member of the DAP-Kinase-related proteins subfamily, was recently identified (Kawai et al., 1998; Kogel et al., 1998). Unlike DAP-Kinase and DRP-1, ZIP-kinase is a nuclear protein, which instead of being regulated by a calmodulin-binding domain, is activated only by homo-dimerization via its leucine-zipper motifs (Kawai et al., 1998). To this group of kinases, another two less homologous nuclear proteins, DRAK1 and DRAK2, were recently added (Sanjo et al., 1998). Together they form a novel subfamily of serine/threonine kinases, as is evident from multiple sequence and phylogenetic analyses (Inbal et al., 1999). A multiple sequence alignment of the kinase domain of these serine/threonine kinases is shown in Fig. 2A.

To check the cellular functions of DRP-1, the laboratory of the present inventor overexpressed wild type DRP-1 in various cell lines and found that it induced apoptosis as measured by a few parameters. Unlike the wild type DRP-1, a kinase inactive mutant of DRP-1 (DRP-1 K42A), did not induce apoptosis, although it was expressed at a similar level in the transfected cells. *In vitro* kinase assays confirmed that DRP-1 K42A is indeed unable to phosphorylate MLC or under autophosphorylation. Also, a truncated form of DRP-1 which lacks the CaM-regulatory region, could be shown to induce very high levels of apoptosis, in a similar way to the analogous truncation of the CaM-regulatory region of DAP-Kinase (Δ CaM; Cohen et al., 1997). Such dependence on the catalytic activity for the apoptotic function is apparent also in the other members of DAP-kinase-related proteins (Kawai et al., 1998; Sanjo et al., 1998).

In a deletion mutant study, which is also presented in the Example herein, the existence of a positive element responsible for apoptotic induction which is located at the C-terminal part of DRP-1 is confirmed. This C-terminal tail of DRP-1 is also essential for its dimerization. Thus, self-dimerization is a requirement for the functionality of this kinase in apoptotic assays, although this property can be overridden by a further deletion of the CaM-regulatory region. Like DRP-1, ZIP-kinase-induced cell death is also controlled by its ability to undergo homo-dimerization via the C-terminal leucine zipper domain (Kawai et al., 1998). Three point mutations in the leucine zipper domain abolished the homo-dimerization as well as

the ability of ZIP-kinase to undergo autophosphorylation *in vitro* and significantly reduced its ability to induce cell death of NIH 3T3 cells. It seems reasonable to assume that activation of these kinases is achieved by homo-dimerization followed by trans-phosphorylation events.

The high homology in the kinase domains of DAP-kinase and DRP-1, and the finding that they are both localized to the cytoplasm (either in soluble or insoluble forms), imply that they may share the same or closely related substrates. The phosphorylation sites for these kinases on the substrate may be either different or identical. Thus, these kinases may cooperate to induce apoptosis in the same cell type or, alternatively, function independently in different cell types, tissues or organs, or in response to different stimuli or time windows.

Another possibility is that these kinases act sequentially along the same signaling pathway to induce apoptosis.

The present invention thus provides for the polypeptide of DRP-1 and for a calmodulin-dependent serine/threonine kinase homologue having the properties of DRP-1, such as the ability to phosphorylate protein in a calcium/calmodulin dependent manner and the ability to induce programmed cell death or apoptosis, and having at least 85% sequence identity to the amino acid sequence SEQ ID NO:2 of DRP-1. Preferably, the calmodulin-dependent serine/threonine kinase homologue has at least 90% sequence identity, and more preferably, at least 95% sequence identity to SEQ ID NO:2.

The term "sequence identity" as used herein means that the amino acid sequences are compared by alignment according to Hanks and Quinn (1991) with a refinement of low homology regions using the Clustal-X program. Such an amino acid alignment is shown in Figs. 2A and 2B where the identical amino acid residues are presented in boxes (cutoff=50%) and homologous amino acid residues, determined according to the PAM 250 matrix, are presented by gray shading (cutoff=65%).

The Clustal-X program referred to in the previous paragraph is the Windows interface for the ClustalW multiple sequence alignment program (Thompson et al., 1994). The Clustal-X program is available over the internet at <ftp://ftp-igbmc.u-strasbg.fr/pub/clustalx/>. Of course, it should be understood that if this link becomes inactive, those of ordinary skill in the art can find versions of this program at other links using standard internet search techniques without undue experimentation. Unless otherwise specified, the most recent version of any program referred herein, as of the effective filing date of the present application, is the one which is used in order to practice the present invention.

If the above method for determining "sequence identity" is considered to be nonenabled for any reason, then one may determine sequence identity by the following technique. The sequences are aligned using Version 9 of the Genetic Computing Group's GDAP (global alignment program), using the default (BLOSUM62) matrix (values -4 to +11) with a gap open penalty of -12 (for the first null of a gap) and a gap extension penalty of -4 (per each additional consecutive null in the gap). After alignment, percentage identity is calculated by expressing the number of matches as a percentage of the number of amino acids in the claimed sequence.

In addition to the full length polypeptide of DRP-1 or a functional homologue thereto with at least 85% sequence identity, the present invention also provides for a fragment of the DRP-1 protein of SEQ ID NO:2 which either maintains the ability to induce cell death or lacks this ability but instead is capable of inhibiting the cell killing ability of DRP-1 protein or its functional homologue described above. It was unexpectedly discovered by the present inventor that the 40 amino acid C-terminal tail (residues 321 to 360 of SEQ ID NO:2) is critical to induction of cell death. As the action of DRP-1 is dependent on dimerization, the 40 amino acid tail, by itself, can inhibit the ability of DRP-1 to induce cell death by interfering with and preventing DRP-1 from dimerizing. Furthermore, it was also unexpectedly discovered that the catalytic domain, by itself (without the calmodulin regulatory domain and the 40 amino acid C-terminal tail, e.g., amino acid residues 13 to 275 of SEQ ID NO:2), is super-killing. One of ordinary skill in the art can readily obtain fragments of the full length sequence of the present invention using N-terminal amino peptidases or C-terminal carboxypeptidases. Each fragment can then be readily tested to see if it possesses one of the two functions described herein for such fragments, without undue experimentation.

Besides fragments of DRP-1 having the above-mentioned properties, fragments having an amino acid sequence with at least 85% sequence identity to the above fragments of DRP-1, preferably with at least 90% sequence identity, and more preferably with at least 95% sequence identity, and maintaining the cell death induction or inhibition properties of the original fragment, are also comprehended by the present invention.

Also comprehended by the present invention are chemical derivatives of the DRP-1 and functional homologues and fragments thereof, as defined above, where a "chemical derivative" contains additional chemical moieties not normally part of the DRP-1 amino acid sequence. Covalent modifications of the amino acid sequence are included within

the scope of this invention. Such modifications may be introduced into DRP-1 or fragments thereof by reacting targeted amino acid residues of the peptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues.

Cysteinyl residues most commonly are reacted with alpha-haloacetates (and
5 corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with bromotrifluoroacetone, alpha-bromo- beta-(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl-2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4- nitrophenol, or chloro-7-nitrobenzo-
10 2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylprocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Parabromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0.

15 Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing alpha-amino acid-containing residues include imidoesters, such as methyl picolinimide, pyridoxal phosphate, pyridoxal, chloroborohydride, trinitrobenzenesulfonic acid, O-methylisourea,
20 2,4-pentanedione, and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3- butanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the
25 groups of lysine, as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues *per se* has been studied extensively, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and e-nitro
30 derivatives, respectively.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides (R'N-C-N-R') such as 1-cyclohexyl-3-[2-morpholinyl-(4-ethyl)]

carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl)carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

5 Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

The present invention also comprehends an isolated DNA molecule which includes a nucleotide sequence encoding the DRP-1 protein of SEQ ID NO:2, a functional
10 homologue thereof as described above, or a fragment of DRP-1 which either maintains the ability of DRP-1 to induce cell death or lacks this ability but is instead capable of inhibiting the cell killing ability of DRP-1 protein, as defined above. The isolated DNA molecule according to the present invention is also intended to comprehend a DNA molecule which hybridizes under moderately stringent, preferably highly stringent, conditions to the nucleotide sequence
15 encoding DRP-1 (corresponding to nucleotides 62 to 1141 of SEQ ID NO:1) and which encodes a polypeptide which maintains the cell death induction properties of DRP-1. The present invention further comprehends isolated DNA molecules which hybridize under moderately stringent, preferably highly stringent, conditions to a nucleotide sequence which encodes for a fragment of DRP-1 which either maintains the ability of DRP-1 to induce cell
20 death (i.e., nucleotides 98 to 886 of SEQ ID NO:1 encoding the catalytic kinase domain of DRP-1) or lacks the ability but is instead capable of inhibiting the cell killing ability of DRP-1 protein (i.e., nucleotides 1022 to 1141 of SEQ ID NO:1 encoding the 40 amino acid C-terminal tail of DRP-1). Furthermore, polypeptides encoded by any nucleic acid, such as DNA or RNA, which hybridizes to the nucleotide sequence of nucleotides 62 to 141 of SEQ ID
25 NO:1 under moderately stringent or highly stringent conditions are considered to be within the scope of the present invention as long as the encoded polypeptide maintains the ability of DRP-1 to induce cell death.

As used herein, stringency conditions are a function of the temperature used in the hybridization experiment, the molarity of the monovalent cations and the percentage of
30 formamide in the hybridization solution. To determine the degree of stringency involved with any given set of conditions, one first uses the equation of Meinkoth et al. (1984) for

determining the stability of hybrids of 100% identity expressed as melting temperature T_m of the DNA-DNA hybrid:

$$T_m = 81.5^{\circ}\text{C} + 16.6 (\text{Log}M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$$

where M is the molarity of monovalent cations, %GC is the percentage of G and C nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. For each 1°C that the T_m is reduced from that calculated for a 100% identity hybrid, the amount of mismatch permitted is increased by about 1%.

Thus, if the T_m used for any given hybridization experiment at the specified salt and formamide concentrations is 10°C below the T_m calculated for a 100% hybrid according to the equation of Meinkoth, hybridization will occur even if there is up to about 10% mismatch.

As used herein, "highly stringent conditions" are those which provide a T_m which is not more than 10°C below the T_m that would exist for a perfect duplex with the target sequence, either as calculated by the above formula or as actually measured.

"Moderately stringent conditions" are those which provide a T_m which is not more than 20°C below the T_m that would exist for a perfect duplex with the target sequence, either as calculated by the above formula or as actually measured. Without limitation, examples of highly stringent ($5\text{-}10^{\circ}\text{C}$ below the calculated or measured T_m of the hybrid) and moderately stringent ($15\text{-}20^{\circ}\text{C}$ below the calculated or measured T_m of the hybrid) conditions use a wash solution of 2 X SSC (standard saline citrate) and 0.5% SDS (sodium dodecyl sulfate) at the appropriate temperature below the calculated T_m of the hybrid. The ultimate stringency of the conditions is primarily due to the washing conditions, particularly if the hybridization conditions used are those which allow less stable hybrids to form along with stable hybrids. The wash conditions at higher stringency then remove the less stable hybrids. A common hybridization condition that can be used with the highly stringent to moderately stringent wash conditions described above is hybridization in a solution of 6 X SSC (or 6 X SSPE (standard seline-phosphate-EDTA)), 5 X Denhardt's reagent, 0.5% SDS, 100 $\mu\text{g/ml}$ denatured, fragmented salmon sperm DNA at a temperature approximately 20° to 25°C below the T_m . If mixed probes are used, it is preferable to use tetramethyl ammonium chloride (TMAC) instead of SSC (Ausubel, 1987, 19989).

Additional aspects of the present invention are vectors which carry the isolated DNA molecule according to the present invention and a host cell which is transformed with the isolated DNA according to the present invention.

The present invention further provides for antisense RNA complementary to at least a portion of a messenger RNA (mRNA or "sense" RNA) molecule which is the transcription product of the DNA sequence encoding the DRP-1 protein of SEQ ID NO:2. The antisense DRP-1 sequence can be chemically synthesized or it can be expressed in host cells. However, when expressed in host cells, the expressed antisense RNA must be stable (i.e., does not undergo rapid degradation). Moreover, the antisense DRP-1 RNA, will essentially specifically only hybridize to the sense DRP-1 mRNA and form a stable double-stranded RNA molecule that is essentially non-translatable. In other words, the antisense DRP-1 RNA prevents the expressed sense DRP-1 mRNA from being translated into active DRP-1 protein. When expressed in host cells, a vector-borne antisense DRP-1 sequence may carry either the entire DRP-1 gene sequence or merely a portion thereof as long as the antisense DRP-1 sequence is capable of hybridizing to "sense" DRP-1 mRNA to prevent its translation into DRP-1 protein. Accordingly, an "antisense" sequence of the present invention can be defined as a sequence which is capable of specifically hybridizing to "sense" DRP-1 mRNA to form a non-translatable double-stranded RNA molecule.

The antisense DRP-1 sequence need not hybridize to the entire length of the DRP-1 mRNA. Instead, it may hybridize to selected regions, such as the 5'-untranslated sequence, the coding sequence, or the 3'-untranslated sequence of the "sense" mRNA. In view of the size of the mammalian genome, the antisense DRP-1 sequence is preferably at least 17, more preferably at least 30, base pairs in length. However, shorter sequences may still be useful, i.e., they either fortuitously do not hybridize to other mammalian sequences, or such "cross-hybridization" does not interfere with the metabolism of the cell in a manner and to a degree which prevents the accomplishment of an object of this invention. The greater the length of the antisense sequence and the greater the number of complementary base pairs, the greater the number of non-complementary bases that can be tolerated, especially if the non-complementary bases are scattered. Both the preferred hybridization target on DRP-1 and the preferred antisense sequence length are readily determined by systematic experiment.

Standard methods such as described in Sambrooke et al., (1989) can be used to systematically remove an increasingly larger portion of the antisense DRP-1 sequence from a plasmid vector. Besides the full length antisense DRP-1 sequence, a series of staggered deletions may be generated, preferably at the 5'-end of the antisense DRP-1 sequence. This creates a set of truncated antisense DRP-1 sequences that still remain complementary to

preferably the 5'-end of the sense DRP-1 mRNA and as a result, still forms a RNA molecule that is double-stranded at the 5'-end of the sense DRP-1 mRNA (complements the 3'-end of an antisense DRP-1 RNA) and remains non-translatable.

The antisense RNA according to the present invention can be used in a method
5 to neutralize a mRNA molecule, which is the transcription product of the DNA sequence encoding the DRP-1 protein of SEQ ID NO:2, by allowing the antisense RNA to hybridize to the DRP-1 mRNA to prevent its translation into DRP-1 protein.

A further aspect of the present invention is directed to a composition, such as a pharmaceutical composition, which contains DRP-1, functional homologues or fragments
10 thereof and a pharmaceutically-acceptable excipient, carrier, diluent, or auxiliary agent.

An antibody, which specifically recognizes DRP-1 or functional homologues thereof is part of the present invention as long as the antibody does not cross-react with DAP-Kinase or ZIP-kinase. For instance, an antibody that specifically recognizes the unique 40 amino acid C-terminal tail of DRP-1, which is not present in DAP-Kinase or ZIP-kinase, is a
15 preferred embodiment of the antibody according to the present invention. Such an antibody can be used for diagnostic imaging, purification of DRP-1 etc.

The terms "antibody" or "antibodies" as used herein are intended to include intact antibodies, such as polyclonal antibodies or monoclonal antibodies (mAbs), as well as proteolytic fragments thereof such as the Fab or F(ab')₂ fragments. Furthermore, the DNA
20 encoding the variable region of the antibody can be inserted into other antibodies to produce chimeric antibodies (see, for example, U.S. Patent 4,816,567) or into T-cell receptors to produce T-cells with the same broad specificity (Eshhar et al., 1990; Gross, et al., 1989). Single chain antibodies can also be produced and used. Single chain antibodies can be single chain composite polypeptides having antigen binding capabilities and comprising a pair of
25 amino acid sequences homologous or analogous to the variable regions of an immunoglobulin light and heavy chain (linked V_H-V_L or single chain F_V). Both V_H and V_L may copy natural monoclonal antibody sequences or one or both of the chains may comprise a CDR-FR construct of the type described in U.S. Patent 5,091,513 (the entire contents of which are hereby incorporated herein by reference). The separate polypeptides analogous to the variable
30 regions of the light and heavy chains are held together by a polypeptide linker. Methods of production of such single chain antibodies, particularly where the DNA encoding the polypeptide structures of the V_H and V_L chains are known, may be accomplished in

accordance with the methods described, for example, in U.S. Patents 4,946,778, 5,091,513 and 5,096,815, the entire contents of each of which are hereby incorporated herein by reference.

As mentioned above, the terms "antibody" or "antibodies" are also meant to include both intact molecules as well as fragments thereof, such as, for example, Fab and $F(ab')_2$, which are capable of binding antigen. Fab and $F(ab')_2$ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl et al., 1983). It will be appreciated that Fab and $F(ab')_2$ and other fragments of the antibodies useful in the present invention may be used for the detection and quantitation of DRP-1 or functional homologues thereof according to the methods used for intact antibody molecules. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce $F(ab')_2$ fragments).

The present invention comprehends not only the intact antibodies or fragments, but also any molecule which includes an antigen binding portion of an antibody such that the molecule is capable of binding to the antigen. It is well within the skill of the art for the artisan to make e.g., fusion proteins which include antigen binding portions of an antibody fused to any other material which is desired to be carried to the antigen binding site, such as marker molecules, toxins, etc.

The antibodies, or fragments of antibodies, of the present invention may be used to quantitatively or qualitatively detect the presence of DRP-1 or functional homologues according to the present invention in a sample. The antibody according to the present invention may also be used for the isolation and purification of DRP-1 or homologues and fragments thereof, such as in an affinity column where the antibodies are immobilized on a solid phase support or carrier.

By "solid phase support or carrier" is intended any support capable of binding antigen or antibodies. Well-known supports, or carriers, include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a

bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

5 One of the ways in which the DRP-1-specific antibody can be detectably labeled is by linking the same to an enzyme and used in an enzyme immunoassay (EIA). This enzyme, in turn, when later exposed to an appropriate substrate, will react with the substrate in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means. The detection can be accomplished by
10 colorimetric methods which employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

 Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it
15 is possible to detect DRP-1 protein through the use of a radioimmunoassay (RIA) (Chard, T., "An Introduction to Radioimmune Assay and Related Techniques" (In: Work, T.S., *et al.*, Laboratory Techniques in Biochemistry in Molecular Biology, North Holland Publishing Company, New York (1978), incorporated by reference herein). The radioactive isotope can be detected by such means as the use of a gamma counter or a liquid scintillation counter or by
20 autoradiography. Radioactively labeled antibodies or antibody fragments can also be used for their capacity to kill cells bound by such antibodies, or cells in the immediate vicinity which are exposed to the radiation from such antibodies. It is also possible to label the antibody with a fluorescent compound, a chemiluminescent or bioluminescent compound.

 The antibody molecules of the present invention may also be adapted for
25 utilization in an immunometric assay (also known as a "two-site" or "sandwich" assay) which is well known in the art.

 In the present specification, the term "programmed cell death" is used to denote a physiological type of cell death which results from activation of some cellular mechanisms, i.e., death which is controlled by the cell's machinery. Programmed cell death
30 may, for example, be the result of activation of the cell machinery by an external trigger, e.g., a cytokine, which leads to cell death. The term "apoptosis" is also used interchangeably with programmed cell death.

The term "tumor" in the present specification denotes an uncontrolled growing mass of abnormal cells. This term includes both primary tumors, which may be benign or malignant, as well as secondary tumors, or metastases, which have spread to other sites in the body.

5 DRP-1 can be used to inhibit growth and metastasis of tumors. Tumor cells are exposed to a variety of death-inducing signals which, in combination with DAP-kinase-related I, can lead to death of the tumor cells. For example, in the blood stream, invading tumor cells must resist programmed cell death that is induced by interactions with cytotoxic T lymphocytes, natural killer cells, and macrophages, and with the cytokines which
10 these hematopoietic cells secrete (e.g., IFNs, TNF, IL-1 β). Tumor cells must also resist the apoptotic cell death induced by nitric oxide anions produced by the endothelial cells, and withstand mechanical shearing forces caused by hemodynamic turbulence. Moreover, during the intravasation or extravasation processes, and during growth in a foreign hostile microenvironment, locally produced inhibitory cytokines (e.g., TGF- β or loss of cell-matrix
15 interactions (e.g., detachment from the basement membranes) also trigger apoptotic cell death.

DRP-1 is useful in promoting death of tumor cells. The protein may be administered to patients, in particular, to cancer patients, which administration may cause death of the tumor cells. The protein may be administered *per se*, or may be administered by an expression vector comprising a DNA molecule of the present invention.

20 Because DRP-1 displays 80% identity with the catalytic domain of DAP-kinase and has a region which displays a high homology to the calmodulin-regulatory region of DAP-kinase, it is expected that DRP-1 has enzymatic kinase activity, which is calmodulin-dependent. Thus, DRP-1 has use as an enzyme and may be used, for example, as the enzyme in any *in vitro* enzymatic reaction which requires the presence of a kinase enzyme.
25 Accordingly, DRP-1 can be used *in vitro* to catalyze phosphorylation reactions as a kinase.

DRP-1 is capable of inducing apoptotic cell death when overexpressed in various cell lines. This ectopic cell death is blocked specifically by the death domain of DAP-kinase, suggesting possible crosstalk between these two kinases. Thus, DRP-1 may also be used for promoting the death of normal or tumor cells and for suppressing the metastatic
30 activity of tumor cells. A particular application of the death-promoting aspect is in therapy of diseases or disorders associated with uncontrolled, pathological cell growth, e.g., cancer (primary tumors and metastasis), psoriasis, autoimmune disease and others. Indeed, it is

expected that the DAP-kinase-related protein I of the present invention and DNA encoding it, may be used in the same manner as disclosed in detail in U.S. applications 08/810,712 and 08/631,097, as well as WO 95/10630.

According to a further aspect of the present invention, referred to herein at times as "the screening aspect", 5
times as "the screening aspect", DRP-1 DNA molecules are used in order to screen individuals for predisposition to cancer. In accordance with this aspect the screening is carried out by comparing the sequence of each of the DAP-kinase-related I DNA molecules to each of the respective DAP genes in the individual, or by following RNA and/or protein expression. The absence of a DAP-kinase-related I gene, a partial deletion or any other difference in the
10 sequence that indicates a mutation in an essential region, or the lack of a DRP-1 RNA and/or protein which may result in a loss of function may lead to a predisposition for cancer. For screening, preferably a battery of related DAP and DRP-1 genes maybe used, as well as different antibodies.

In the screening aspect, DAP-kinase related product I molecules may also be
15 used for prognostic purposes. For example, if a tumor cell lacks DRP-1 activity, this may reflect high chances of developing metastasis. In addition, DRP-1 positive cells may be more susceptible to control by chemotherapeutic drugs that work by inducing apoptosis, so that the choice of treatment modalities may be made based upon the DRP-1 state of the cells.

The DAP-kinase-related product can be used to screen individuals for
20 predisposition to cancer. There is provided a method for detecting the absence of a DRP-1 gene, a partial deletion or a mutation (i.e., point mutation, deletion or any other mutation) in the DRP-1 genes of an individual, or the absence of a DRP-1 RNA or protein, comprising probing genomic DNA, cDNA, or RNA from the individual with a DNA probe or a multitude of DNA probes having a complete or partial sequence of the DRP-1 genes, or probing protein
25 extracts with specific antibodies.

A particular application of the screening aspect of this invention is in the screening for individuals having a predisposition to cancer, an absence of the gene, or a detected mutation or deletion indicating that the individual has such a predisposition.

One example of a method in accordance with the screening aspect typically
30 comprises the following steps:

(a) obtaining a sample of either genomic DNA from cells of the individual or cDNA produced from mRNA of said cells;

(b) adding one or more DNA probes, each of said probes comprising a complete or partial sequence of a DRP-1 gene;

(c) providing conditions for hybridization to determine whether the DRP-1 gene is present or absent, i.e., whether there is a match between the sequence of the DNA probe or probes and a sequence in the DNA of said sample or a mismatch, a mismatch indicating a deletion or a mutation in the endogenous DNA and a predisposition to cancer in the tested individual.

Other examples of the screening aspect of the invention are well known to the skilled artisan and include, but are not limited to, Northern blots, RNase protection assays, and various PCR procedures.

The mutation in the DRP-1 gene, indicating a possible predisposition to cancer, can also be detected by the aid of appropriate antibodies which are able to distinguish between a mutated, a non-functional and a normal functional DRP-1 gene product. In addition, mutations that abolish protein translation or transcription due to promoter inactivation can be detected with the aid of antibodies that are used to react with protein cell extracts. Screening is also possible with respect to metastases.

Having now generally described the invention, the same will be more readily understood through reference to the following example which is provided by way of illustration and is not intended to be limiting of the present invention.

EXAMPLE

In this study, the identification and the structure/function analysis of a novel DAP-kinase-related protein, DRP-1, is described, DRP-1 is a 42kDa Ca^{2+} /CaM-regulated serine/threonine kinase which shows high degree of homology to DAP (Death Associated Protein)-kinase. The homology spans over the catalytic domain and the calmodulin-regulatory region, whereas the rest C-terminal part of the protein differs completely from DAP-kinase and displays no homology to any known protein. The catalytic domain is also homologous to the recently identified ZIP-kinase and to a lesser extent to the catalytic domains of DRAK1/2, thus forming together a novel subfamily of serine/threonine kinases. DRP-1 is localized to the cytoplasm as shown by immunostaining and cellular fractionation assays. *In vitro* kinase assays indicate that wild type DRP-1, but not a kinase inactive mutant, undergoes autophosphorylation and phosphorylates an external substrate in a Ca^{2+} /CaM-dependent

manner. Ectopically expressed DRP-1 is able to induce apoptosis in various types of cells. Cell killing by DRP-1 is dependent on two features: the intact kinase activity and the presence of C-terminal 40 amino acids shown to be involved in self-dimerization of the kinase. Interestingly, further deletion of the CaM-regulatory region overrode the indispensable role of the C-terminal tail and generated a "super-killer" mutant. Finally, a dominant negative fragment of DAP-kinase encompassing the death domain is a potent blocker of apoptosis induced by DRP-1. This implies a possible functional connection between DAP-kinase and DRP-1. The experiments conducted in this study and the results obtained are presented below.

MATERIALS AND METHODS

cDNA cloning and Northern Blot Analysis

A PCR fragment of 364 bp was obtained from a λ gt11 human spleen cDNA library (Clontech) using primers from the deduced DRP-1 sequence, 1047-GGCCGGATGAGGACCTGAGG-1066 (SEQ ID NO:13) and 1411-TCCACACTCCCACCCCAGACTC-1390 (SEQ ID NO:14). To obtain the full length cDNA of DRP-1, the same cDNA library was screened with the radiolabeled PCR product. Positive phage clones were isolated, cDNA was subcloned into a BlueScript vector and analyzed by restriction enzyme mapping and DNA sequencing. A 270 bp 3'-fragment from the full length cDNA of DRP-1 was generated by EcoRI digestion, and used to probe polyA+ RNA prepared by a standard procedure from various cell lines.

In vitro Transcription and Translation Assay

The full length cDNA was used as a template for *in vitro* transcription from the T7 promoter. This RNA was translated in reticulocyte lysate (TNT® T7 Quick Coupled Transcription/Translation System; Promega) by conventional procedures, with [35 S] methionine (Amersham) as a labeled precursor. The reaction product was then run on 12% SDS-PAGE gel, followed by sodium salicylate incubation for signal amplification. The gel was dried and exposed to X-ray film.

In vitro Kinase Assay

293 cells were transfected by a FLAG-tagged wild type DRP-1, DRP-1 K42A mutant, or mock-transfected. Cell lysates of 293 transfected cells were prepared as previously described (Deiss et al., 1995). Immunoprecipitation of DRP-1 or DRP-1 K42A mutant from 150µg total extract was done with 20µl anti-FLAG M2 gel (IBI, Kodak) in 500µl of PLB supplemented with protease and phosphatase inhibitors for 2h at 4°C. Following three washes with PLB, the immunoprecipitates were washed once with reaction buffer (50 mM HEPES pH 7.5, 20 mM MgCl₂, and 0.1 mg/ml BSA). The proteins bound to the beads were incubated for 15 min at 30°C in 50 µl of reaction buffer containing 15 µCi [γ 32p] ATP (3 pmole), 50µM ATP, 5µg MLC (Sigma), and where indicated, also 1µM bovine calmodulin (Sigma), 0.5 mM CaCl₂, or 3mM EGTA in the absence of calmodulin/CaCl₂. Protein sample buffer was added to terminate the reaction, and after boiling, the proteins were analyzed on 11% SDS-PAGE. The gel was blotted onto a nitrocellulose membrane and ³²P- labeled proteins were visualized by autoradiography.

15 Immunostaining of Cells

DRP-1 transfected or mock-transfected COS-7 cells were plated on glass cover-slips (13 mm diam.). After 48 hours, the cells were fixed/permeabilized in 3% formaldehyde for 5 min, methanol 5 min, acetone 2 min. The cells were blocked in 10% NGS for 30 min and incubated with anti-FLAG antibodies (dilution 1:100; IBI, Kodak) in 10% NGS for 60 min. Rhodamine-conjugated goat anti-mouse secondary antibodies (dilution 1:200; Jackson Immuno Research Lab.) and the nucleic acid dye, Oligreen (dilution 1:5000; Molecular Probes), for nuclear staining were then applied. The coverslips were mounted in Mowiol and observed under fluorescence microscope.

25 Detergent Extraction Assay

Sub-confluent cultures of COS-7 transfected cells, grown on 9 cm plate, were washed once with PBS and then with MES buffer (50 mM MES pH 6.8, 2.5 mM EGTA, 2.5 mM MgCl₂). The cells were extracted for 3 min with 0.5 ml of 0.5% Triton X-100 in MES buffer supplemented with protease inhibitors. The supernatant (the soluble fraction- Sol) was collected, centrifuged for 2 min. at 16,000x g at 4°C, and the clear supernatant was then transferred to new tubes. Two volumes of cold ethanol were added and the tubes were incubated at -20°C for overnight, centrifuged 10 min. at 16,000x g at 4°C and resuspended in

200 μ l of 2x protein sample buffer without dye. The detergent insoluble matrix (InSol) remaining on the plate was extracted in 200 μ l of 2x protein sample buffer, scraped from the plate with rubber policeman and collected into tube. The samples were loaded on 10% SDS-PAGE, 100 μ g protein extracts were loaded on each lane from the Sol fraction,

- 5 equivalent volumes of InSol were loaded. Analysis of the proteins was done using monoclonal anti-FLAG antibodies (dilution 1:200; IBI, Kodak).

Cell Lines, Transfections and Apoptotic Assays

- All cell lines were grown in DMEM (Biological Industries) supplemented with
10 10% fetal calf serum (Bio-Lab). For transient transfection, 1×10^5 cells per well, were seeded in a 6 well plate a day before transfection. Transfections were done by calcium-phosphate method. For cell death assays by inducing overexpression, a mixture containing 1.5 μ g of cell death plasmid (expressing either DRP-1 or Δ CaM DAPk mutant) and 0.5 μ g of pEGFP-NI plasmid (Clontech) was used. For cell death protection assays we used a mixture containing
15 1.2 μ g of cell death inducing plasmid (either DRP-1 or Δ CaM DAPk mutant), 0.5 μ g of a plasmid to be tested for cell death protection (expressing DAPk-DD, DN FADD or luciferase as negative control), and 0.5 μ g of pEGFP-NI plasmid. Cells were counted and photographed 24 hours after transfection. In each transfection, three fields, each consisting of at least 100 GFP-positive cells, were scored for apoptotic cells according to their morphology. When
20 indicated, cell lysates were prepared from the transient transfection at 24 hours, for protein analysis. The transfections of Rat embryo fibroblasts (REF) and FACS analysis of transfected fibroblasts for DNA content distribution were done as previously described (Kissil et al., 1998).

25 Co-immunoprecipitation assays

- 293 cells grown in 90mm plates (1×10^6 cells/plate) were co-transfected with 5 μ g FLAG-tagged or HA-tagged DRP-1 and 20 μ g of HA-tagged or FLAG-tagged RFX1- Δ SmaI, respectively, or with DRP-1-HA and DRP-1-FLAG, 5 μ g each. Immunoprecipitation of DRP-1 or RFX1- Δ SmaI from 1mg total extract was done using anti-FLAG M2 gel or anti-
30 HA as described above. Detection of bound proteins was done using anti-HA antibodies (dilution 1:1000, Babco) or anti-FLAG antibodies. For the deletion mutant study, 5 μ g of FLAG-tagged fully length DRP-1 were co-transfected with 5 μ g of HA-tagged DRP-1 deletion

mutants. Immunoprecipitation of DRP-1 from 1mg total extract was done using anti-FLAG M2 gel as described above. Detection of co-immunoprecipitated proteins (the mutants of DRP-1 or full length DRP-1) was done using anti-HA antibodies.

5 Nucleotide sequence accession number

The nucleotide sequence of human DRP-1 has been submitted to the GenBank™/EBI Data Bank (accession no. AF052941). The murine DRP-1 is also deposited at the GenBank™/EBI Data Bank (accession no. AF052942).

10 RESULTS

Cloning of DRP-1

To identify proteins that share homologous sequences with DAP-kinase, EST databases were searched using the BLAST™ program. Two ESTs of human and murine origin showed remarkable amino acid homology to the catalytic domains of DAP-kinase and the recently identified protein ZIP-kinase (79.5% and 80.2% identity, respectively). A second EST search was performed using the 5' and the 3' ends of the human EST, which identified a few more overlapping ESTs. A putative novel cDNA sequence was generated and used to design primers for cloning the full length cDNA. PCR performed on human spleen cDNA library amplified a 364 bp fragment that was further used to screen the same library. The full length cDNA was then isolated, subcloned into BlueScript vector, and sequenced.

The isolated cDNA was found to be 1742 bp long and to contain a serine/threonine kinase domain with all of the 12 characterized subdomains present (Park et al., 1997, Fig. 1A). Sequence alignment indicated that the catalytic domain of DRP-1 has 80% sequence identity to that of DAP-kinase and ZIP-kinase, yet less 50% sequence identity to the newly identified DRAK proteins (Fig. 2A). Like DAP-kinase but unlike ZIP-kinase, DRP-1 carries a typical CaM-regulatory region adjacent to its catalytic domain, as shown in Figures 1 and 2B. As compared with other kinases such as CaKIIa and MLCK, DRP-1 has the highest homology to DAP- kinase in this region, as shown in Figure 2B. The remaining short stretch of amino acids at the C-terminal part of DRP-1 (40 amino acid tail) displays no homology to any known protein.

30 Expression of DRP-1 and Tissue Distribution

To check the RNA expression of DRP-1, polyA+RNA was prepared from various cell lines and hybridized to a probe designed from the less conserved region of DRP-1. A single weak band of 1.9 kb appeared in some cell lines, in a Northern blot analysis of poly A+RNA (3 micrograms) extracted from various cell lines (Fig. 3A), suggesting that the mRNA is expressed at low amounts in HeLa, 293 and MCF-7 cells. The mRNA was hybridized with a radiolabeled human DRP-1 probe. The position of the transcript is indicated by an arrow. From PCR analysis of various cDNA libraries and the data gathered from EST searches, it was concluded that human DRP-1 is expressed, at least, in spleen, colon, breast, and leukocyte tissue.

In vitro transcription and translation assays conducted in reticulocyte lysates using the cloned DRP-1 cDNA as a template generated a single protein band of about 42 kDa in size, as predicted by its sequence. This protein band, obtained by Western blot analysis of *in vitro* transcribed and then translated DRP-1, is shown in Figure 3B. A FLAG-tagged DRP-1 was then cloned into pCDNA3 vector and expressed in HeLa cells. A protein of 42 kDa was evident upon immunoblot analysis of the cell lysates with anti-FLAG antibodies, shown in Figure 3C. In this case, 24 hours following transfection, the cells were harvested and lysed. The extracted proteins were separated by SDS-PAGE and then immunoblotted with anti-FLAG antibodies.

Cellular Localization of Ectopically Expressed DRP-1

In order to follow the cellular localization of the exogenous DRP-1, FLAG-tagged DRP-1 was expressed in COS-7 cells. COS-7 cells were transfected by a FLAG-tagged DRP-1 cloned in pCDNA3 vector, fixed and permeabilized in 1% formaldehyde followed by methanol/acetone treatment. Cells were visualized under fluorescence microscope. Immunoblot analysis proved that DRP-1 was expressed in these cells. For the immunostaining procedure, the non-transfected (Fig. 4A) and DRP-1 transfected (Fig. 4B) COS-7 cells were then fixed and reacted both with Oligreen for nuclear staining and anti-FLAG antibodies for DRP-1 staining. Specific DRP-1 staining was detected in the cytoplasm of these cells, as shown in Figure 4B.

A gentle cell extraction was performed with nonionic detergent, 0.5% TRITON X-100, that removes lipids and soluble proteins, leaving intact the detergent insoluble matrix composed of the nucleus, the cytoskeleton framework, and cytoskeleton-

associated proteins. In contrast to DAP-kinase, which is exclusively localized to the cytoskeleton, and hence found only in detergent insoluble fractions (Cohen et al., 1997, Fig. 4C), DRP-1 was preferentially eluted from the detergent soluble fraction, while a small amount was eluted from the insoluble fraction, as shown in Figure 4C. Thus, it was concluded that DRP-1 is a cytoplasmic protein with minor association with insoluble matrix components.

Intrinsic Kinase Activity of DRP-1

To test whether DRP-1 functions as a kinase as predicted from the amino acid sequence, an *in vitro* kinase assay was performed using myosin light chain (MLC) as an exogenous substrate. This substrate was chosen because it is phosphorylated by DAP-kinase (Cohen et al., 1997). DRP-1 was transfected into human kidney 293 cells, immunoprecipitated, and incubated with MLC in the presence and absence of Ca^{2+} and calmodulin. Both MLC phosphorylation and DRP-1 autophosphorylation were evident, as can be seen from Figure 5A.

In assaying the *in vitro* kinase activity of DRP-1, the proteins were assayed in the presence or absence of Ca^{2+} /CaM and MLC. The proteins were run on 11% SDS-PAGE and blotted to nitrocellulose membrane. Figure 5A shows the autophosphorylation of DRP-1 and MLC phosphorylation, respectively, as seen after exposure of X-ray film. Figure 5B shows the DRP-1 proteins by incubation of the same blot with anti-FLAG antibodies and ECL detection.

The addition of Ca^{2+} /calmodulin to the reaction mixture increased the amount of phosphorylated MLC, in accordance with the assumption that, like DAP-kinase, DRP-1 is negatively regulated by the autoinhibitory calmodulin binding domain, and that this inhibition is removed by the binding of Ca^{2+} /calmodulin. A catalytically inactive mutant of DRP-1, DRP-1 K42A, did not phosphorylate MLC and failed to undergo autophosphorylation even though higher amounts of DRP-1 protein were present, as can be seen from Figure 5A. Thus, DRP-1 was found to function *in vitro* as a kinase that is capable of phosphorylating itself and an external substrate. This latter property is stimulated by the addition of Ca^{2+} and calmodulin.

DRP-1 Induces Apoptosis in a Variety of Cell Lines

The high homology to DAP-kinase in the kinase and calmodulin-binding regions suggested the value of checking whether DRP-1 is involved in apoptosis. The wild type DRP-1 and the catalytically inactive mutant of DRP-1, DRP-1 K42A, which are cloned in pCDNA3 vector, were transfected into 293 cells. To quantitate the number of apoptotic cells, these constructs were transfected with a vector expressing the GFP protein. The GFP protein was used as a marker to visualize the transfected cells and to assess the apoptotic frequency among the transfectants according to morphological alterations. Apoptotic cells were scored after 24 hours. Overexpression of the DRP-1 resulted in massive apoptotic cell death (50-60%), as compared to the basal level of apoptotic cells caused by transfection of the non-relevant gene luciferase, shown in Figures 6A-6B and 7.

Most of the GFP positive green cells rounded up and shrunk; some of them showed cytoplasmic blebs, and some were further fragmented into "apoptotic bodies." In addition, some of the transfected cells detached from the plate. This apoptotic cell death was only slightly lower than that of an activated DAP- kinase mutant lacking the autoinhibitory calmodulin regulatory region (Δ CaM; apoptotic values of 70-80%). In contrast, when the cells were transfected with the kinase inactive mutant of DRP-1, DRP-1 K42A, as shown in Figures 6A-6D and 7, no apoptosis was observed. This experiment was repeated six times with reproducible results.

Western blot analysis of transfected cells, using anti- FLAG antibodies, confirmed the expression of both the exogenous wild type and K42A mutant of DRP-1 (Figures 8A and 8B). Similar results were also observed in human SV-80 fibroblasts. In another type of assay, the effect of ectopically expressed DRP-1 on the DNA content of rat embryo primary fibroblasts (REF cells) was assessed, as previously described (Kissil et al., 1999). The REFS were co-transfected with DRP-1 and a membrane-bound form of GFP and then after 48 hours subjected to FACS analysis of their DNA content. A fraction of cells displaying a sub-G1 population, indicative of cells containing fragmented DNA, appeared exclusively in the DRP-1 transfected cells but not in cells transfected with a control vector or with the DRP-1 K42A mutant form. No effect was found on cell cycle distribution of the viable cells.

To obtain the results shown in Figures 6A-6D, 1×10^5 293 cells/well were co-transfected with FLAG-tagged wild type DRP-1 or K42A mutant of DRP-1, 1.5 microgram/well and GFP, 0.5 microgram/well. GFP positive cells were visualized under

fluorescent microscope and scored for the appearance of apoptotic morphology 24 hours after transfection. Apoptotic cells are indicated by arrows. The fluorescent microscopic images correspond to 293 cells transfected by pCDNA3-luciferase as negative control (Fig. 6A), pCDNA#-deltaCaM DAP-kinase as positive control (Fig. 6B), pCDNA3-DRP-1 (Fig. 6C),
5 pCDNA3-K42A DRP-1 (Fig. 6D).

In Figure 7, graphs show the percentage of apoptotic cells resulting from the above-mentioned transfections (average \pm S.D. calculated from triplicates of 100 cells each). The scores were taken from the same experiment shown in Figures 6A-6D.

10 In Figures 8A and 8B, proteins extracted from the transfected cells were separated on 10% SDS-PAGE and blotted to nitrocellulose membrane. The blot was hybridized with anti-FLAG antibodies for DRP-1 detection and anti-vinculin antibodies to quantitate the loaded protein amounts. The proteins were prepared from the same experiments shown in Figures 6A-6D.

15 DAP Kinase Death Domain Protects From DRP-1 Induced Apoptosis

The structural homology of DRP-1 to DAP-kinase, the common regulation by Ca^{2+} /calmodulin, and the finding that both proteins caused apoptosis upon overexpression, suggested that they function along a common apoptotic pathway. In order to test this possibility, the effect of the dominant-negative DAP- kinase death domain (DAPk DD) on
20 DRP-1-induced cell death was analyzed. The laboratory of the present inventor showed recently that overexpression of the fragment encompassing the death domain of DAP-kinase acts as a specific dominant-negative mutant, negating the effects of the full length protein (Datta et al., 1997). As a consequence, it protected cells from TNF-alpha, Fas and FADD/MORT1-induced cell death (Datta et al., 1997).

25 It has now been discovered that DAPk DD protected cell death induced by DRP-1 in 293 cells. As shown in Figure 9A, the apoptotic ratio dropped from 64.3% to 24.7%. A control transfection including DRP-1 and a non-relevant luciferase DNA excluded the possibility that this blockage was simply due to larger amount of DNA used in the transfection. Moreover, the effect of DAPk DD was specific, since the death domain of
30 FADD failed to manifest a similar effect. (Figure 9A). Western blot analysis of transfected cells using anti-FLAG antibodies confirmed the expression of the exogenous DRP-1 in all transfections, as shown in Figure 9B. This experiment was repeated three times with

reproducible results. The ability of the death domain of DAP- kinase to block death induced by DRP-1 implies that DAP-kinase and DRP-1 function along a common pathway.

To obtain the results shown in Figure 9A, 1×10^5 cells/well of 293 cells were co-transfected with 1.2 microgram/well of FLAG-tagged wildtype DRP-1 and 0.5 microgram/well of GFP. The scores are the percentage of apoptotic cells given as average \pm S.D. and calculated from triplicates of 100 cells each.

To demonstrate the DRP-1 protein expression in 293 transfected cells shown in Figure 9B, proteins extracted from the transfected cells were separated on 10% SDS-PAGE and blotted to nitrocellulose membrane. The blot was hybridized with anti-FLAG antibodies for DRP-1 detection and anti-vinculin antibodies to quantitate the loaded protein amounts. The proteins were prepared from the same experiment shown in Figure 9A.

Deletion of the C-terminal tail of DRP-1 abolishes its apoptotic activity, while further truncation of the CaM-regulatory region strongly enhances the apoptotic effect

In order to further understand the mode of DRP-1 action in apoptosis, constructs containing C-terminal truncations of DRP-1 tagged by HA were constructed (Fig. 10A). DRP-1 $\Delta 40$ lacks the most C-terminal part of DRP-1 which displays no homology to any known protein. DRP-1 $\Delta 73$ lacks, in addition to that, the CaM-regulatory region of DRP-1, and DRP-1 $\Delta 85$ contains only the catalytic domain. The wild type DRP-1 and the various truncation mutants of DRP-1 were transfected into 293 cells. Induction of apoptotic cell death was assayed as mentioned above in DRP-1 induced apoptosis. Overexpression of the wild type DRP-1 resulted in apoptosis (25%) while the DRP-1 $\Delta 40$ had no effect in these assays. On the other hand, further truncations of the CaM-regulatory region, yielded mutants ($\Delta 73$, $\Delta 85$) which acted as "super-killers" (~90% apoptosis) (Figs. 11A and 11B). This experiment was repeated three times with reproducible results. Western blot analysis of transfected cells, using anti-HA antibodies confirmed the expression of all DRP-1 forms (Fig. 10B). Thus, the dependence of the apoptotic effect of DRP-1 on its kinase activity was confirmed again, since removal of the CaM-regulatory region which acts as an autoinhibitory domain generates a constitutively active kinase. In addition, the existence of a positive module in the C-terminal region of DRP-1, which is necessary for its pro-apoptotic effect, provided that the CaM-regulatory effect is still present, is shown.

The C-terminal part of DRP-1 functions as a homo-dimerization domain

Western analysis performed on proteins extracted from 293 cells transfected by FLAG-tagged DRP-1 revealed an additional band (not shown). This observation led the present inventor to test whether DRP-1 can undergo homo-dimerization. To this aim, two constructs expressing DRP-1 fused to either FLAG or HA tags were co-transfected into 293 cells and classical pull-down experiments with each one of the two epitopes were performed. FLAG-tagged DRP-1 could be shown to bind specifically to HA-tagged DRP-1 in both IP directions (Fig. 12A, see lane 3 in both IP Panels). No binding of DRP-1-HA to FLAG beads or to the irrelevant cytoplasmic protein RFX- Δ SmaI could be observed (Fig. 12A, see IP anti-FLAG panel, lanes 2 or 1+2, respectively). Also non-specific binding of DRP-1-FLAG to HA bead or to RFX- Δ SmaI protein could not be detected (Fig. 12A, see IP anti-HA panel, lanes 1 or 1+2, respectively). Western analyses confirmed the expression of all proteins in these cell extracts (Fig. 12A, see Western panels).

The observation that a C-terminal truncation of 40 amino acids in DRP-1 abolished its apoptotic effect upon ectopic expression in 293 cells, prompted the present inventor to test whether this domain may be involved in the homo-dimerization of DRP-1. DRP-1-FLAG was co-expressed in conjugation with the various deletion mutants of DRP-1 tagged by HA. A strong binding of DRP-1-FLAG to the wild type DRP-1-HA was detected, whereas the binding to DRP-1 Δ 40 was mostly abolished (Fig. 12B, upper IP panel, compare lane 1 to 2-4). Western analysis confirmed the expression of wild type DRP-1-HA and all other DRP-1-HA deletion mutants in these transfections (Fig. 12B, see Western panel). Lower IP panel confirmed the expression of wild type DRP-1-FLAG in all these transfections. Thus, the present inventor concluded that a region spanning the C-terminal 40 amino acids of DRP-1 is responsible for its homo-dimerization. This homo-dimerization is probably required for the apoptotic effect of DRP-1, since DRP-1- Δ 40 has lost the ability to induce apoptosis in 293 cells (Figs. 11A and 11B).

Having now fully described this invention, it will be appreciated by those skilled in the art that the same can be performed within a wide range of equivalent parameters, concentrations, and conditions without departing from the spirit and scope of the invention and without undue experimentation.

While this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This

application is intended to cover any variations, uses or adaptations of the inventions following, in general, the principles of the invention and including such departures from the present disclosure as come within know or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth as follows in the scope of the appended claims.

All references cited herein, including journal articles or abstracts, published or unpublished U.S. or foreign patent application, issued U.S. or foreign patents, or any other references, are entirely incorporated by reference herein, including all data, tables, figures, and text presented in the cited references. Additionally, the entire contents of the references cited within the references cited herein are also entirely incorporated by reference.

Reference to known method steps, conventional method steps, known methods or conventional methods is not in any way an admission that any aspect, description or embodiment of the present invention is disclosed, taught or suggested in the relevant art.

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art (including the contents of the references cited herein), readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance presented herein, in combination with the knowledge of one of ordinary skill in the art.

REFERENCES

- Anderson, P., "Kinase cascades regulating entry into apoptosis," Microbiol. Mo. Biol. Rev., **61**, 33-46, 1997.
- Ausubel et al., Current Protocols in Molecular Biology, Green Publications and Wiley
5 Interscience, New York, 1987-1999.
- Basu, S., and Kolesnick, R., "Stress signals for apoptosis: ceramide and c-Jun kinase,"
Oncogene, **17**, 3277-3285, 1998.
- Bokoch, G.M., "Caspase-mediated activation of PAK2 during apoptosis: proteolytic kinase
activation as a general mechanism of apoptotic signal transduction?," Cell death diff.,
10 **5**, 637-645, 1998.
- Cardone, M.H., Salveson, G.S., Widmann, C., Johnson, G. And Frisch, S.M., "The
regulation of anoikis: MEKK-1 activation requires cleavage by caspases.", Cell,
90, 315-323, 1997.
- Cardone, M.H., Roy, N., Stennicke, H.R., Salvesen, G.S., Franke, T.F., Stanbridge, E.,
15 Frisch, S., and Reed, J.S., "Regulation of cell death protease caspase-9 by
phosphorylation.", Science, **282**, 318-321, 1998.
- Cohen, O., Feinstein, E., and Kimchi, A., "DAP-kinase is a Ca²⁺/calmodulin-dependent,
cytoskeletal-associated protein kinase, with cell death-inducing functions that depend
on its catalytic activity.", EMBO J., **16**, 998-1008, 1997.
- 20 Cohen, O., Inbal, B., Kissil, J.L., Feinstein, E., Spivak, T., and Kimchi, A., "DAP-kinase
participates in TNF- α and Fas-induced apoptosis and its function requires the death
domain.", J. Cell. Biol., in press, 1999.
- Datta, S.R., Dudek, H., Tao, X., Masters, S., Fu, H., Gotoh, Y., and Greenberg, M.E., "Akt
phosphorylation of BAD couples survival signals to the cell-intrinsic death
25 machinery.", Cell, **91**, 231-241, 1997.
- Deiss, L.P., Feinstein, E., Berissi, H., Cohen, O., and Kimchi, A., "Identification of a novel
serine/threonine kinase and a novel 15-kD protein as potential mediators of the
gamma interferon-induced cell death.", Genes Dev., **9**, 15-30, 1995.
- Deiss, L.P. and Kimchi, A., "A genetic tool used to identify thioredoxin as a mediator of a
30 growth inhibitory signal.", Science, **252**, 117-120, 1991.

- del Peso, L., Gonzalez-Garcia, M., Page, C., Herrera, R., and Nunez, G., "Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt.", Science, **282**, 318-321, 1997.
- Eshhar, Z. et al., Br. J. Cancer Suppl., **10**, 27-9, 1990.
- 5 Feinstein, E., Kimchi, A., Wallach, D., Boldin, M., and Varfolomeev, E., "The death domain: a module shared by proteins with diverse cellular functions.", Trends Biochem. Sci., **20**, 342-344, 1995.
- Green, D., and Kroemer, G., "The central executioners of apoptosis: caspases or mitochondria?" Trends Cell Biol., **8**, 267-271, 1998.
- 10 Gross, G. et al., Proc. Natl. Acad. Sci. USA, **86**, 10024-8, 1989.
- Hanks, S.K., and Quinn, A.M., "Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members.", Methods Enzymol., **200**, 38-62, 1991.
- Inbal, B., Cohen, O., Polak-Charcon, S., Kopolovic, J., Vadai, E., Eisenbach, L., and Kimchi, A., "DAP kinase links the control of apoptosis to metastasis.", Nature, **390**, 180-184, 1997.
- 15 Inbal, B., Kissil, J.K., Cohen, O., Spivak-Kroizman, T., and Kimchi, A., "The DAP-related protein kinases-a novel subfamily of serine/threonine kinases with a possible link to apoptosis.", submitted, 1999.
- 20 Jacobson, M.D., Weil, M., and Raff, M.C., "Programmed cell death in animal development." Cell, **88**, 347-354, 1997.
- Kawai, T., Matsumoto, M., Takeda, K., Sanjo, H., and Akira, S., "ZIP kinase, a novel serine/threonine kinase which mediates apoptosis.", Mol. Cell Biol., **18**, 1642-1651, 1998.
- Kelliher, M.A., Grimm, S., Ishida, Y., Kuo, F., Stanger, B.Z., and Leder, P., "The death domain kinase RIP mediates the TNF-induced NF-kappaB signal.", Immunity, **8**, 297-303, 1998.
- 25 Kimchi, A., J. Cell. Biochem., **50**, 1-9, 1992.
- Kimchi, A., "DAP genes: novel apoptotic genes isolated by a functional approach to gene cloning.", Biochim. Biophys. Acta, **1377**, F13-33, 1998.
- 30 Kissil, J.L., and Kimchi, A., "Death-associated proteins: from gene identification to a the analysis of their apoptotic and tumor suppressive functions.", Mol. Med. Today, **4**, 268-74, 1998.

- Kissil, J.L., Cohen, O., Raveh, T., and Kimchi, A., "DAP-kinase loss of expression in various carcinoma and B-cell lymphoma cell lines: possible implications for role as tumor suppressor gene.", EMBO J., **18**, 353-362, 1999.
- Kogel, D., Plottner, O., Landsberg, G., Christian, S., and Scheidtmann, K.H., "Cloning and
5 characterization of Dlk, a novel serine/threonine kinase that is tightly associated with chromatin and phosphorylates core histones.", Oncogene, **17**, 2645-2654, 1998.
- Levy et al., Mol. Cell. Biol., **13**, 7942-7952, 1993.
- Levy-Strumpf, N., and Kimchi, A., "Death associated proteins (DAPs): from gene
10 identification to the analysis of their apoptotic and tumor suppressive functions."
Oncogene, **17**, 3331-3340, 1998.
- Maundrell, K., Antonsson, B., Magnenat, E., Camps, M., Muda, M., Chabert, C., Gillieron, C., Boschert, U., Vial-Knecht, E., Martinou, J.C., and Artkinstall, S., "Bcl-2 undergoes phosphorylation by c-Jun N-terminal kinase/stress-activated protein
15 kinases in the presence of the constitutively active GTP-binding protein Rac1.", J. Biol. Chem., **272**, 25238-25342, 1997.
- McCarthy, J.V., Ni, J., and Dixit, V.M., "RIP2 is a novel NF-kappaB-activating and cell death-inducing kinase, J. Biol. Chem., **273**, 16968-75, 1998.
- Meinkoth et al., Anal. Biochem., **138**, 267-284, 1984.
- Park, J., Kim, I., Oh, Y.J., Lee, K., Han, P.L., and Choi, E.J., "Activation of c-Jun N-
20 terminal kinase antagonizes an anti-apoptotic action of Bcl-2.", J. Biol. Chem., **272**, 16725-16728, 1997.
- Peitenpol et al., Cell, **61**, 777-785, 1990.
- Sambrooke et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY, 1989.
- 25 Sanjo, H., Kawai, T., and Akira, S., "DRAKS, novel serine/threonine kinases related to death-associated protein kinase that trigger apoptosis.", J. Biol. Chem., **273**, 29066-29071, 1998.
- Stanger, B.Z., Leder, P., Lee, T.H., Kim, E., and Seed, B., "RIP: a novel protein containing a death domain that interacts with Fas/APO-1 (CD95) in yeast and causes cell
30 death.", Cell, **81**, 513-523, 1995.
- Sun, X., Lee, J., Navas, T., Baldwin, D.T., Stewart, T.A., and Dixit, V.M., "RIP3, a Novel Apoptosis-inducing Kinase.", J. Biol. Chem., **274**, 16871-16875, 1999.

- Thompson, J.D., Higgins, D.G., Gibson, T., J., "CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice.", Nucleic Acids Res., **22**, 4673-4680, 1994.
- 5 Verheij, M., Ruiter, G.A., Zerp, S.F., Van Blitterswijk, W.J., Fuks, Z. Haimovitz-Friedman, A., and Bartelink, H., "The role of the stress-activated protein kinase (SAPK/JNK) signaling pathway in radiation-induced apoptosis.", Radiother. Oncol., **47**, 225-232, 1998.
- Wahl et al., J. Nucl. Med., **24**, 316-325, 1983.
- 10 White, E., "Life, death and the pursuit of apoptosis.", Genes Dev., **10**, 1-15, 1996.
- Yang, X., Khosravi-Far, R., Chang, H.Y., and Baltimore, D., Cell, **89**, 1067-1076, 1997.
- Yu, P.W., Huang, B.C., Shen, M., Quast, J., Chan, E., Xu, X, Nolan, G.P., Payan, D.G. and Luo, Y., "Identification of RIP3, a RIP-like kinase that activates apoptosis and Nf-kappa B.", Curr. Biol., **9**, 539-42, 1999.

WHAT IS CLAIMED IS:

1. An isolated polypeptide, which is a calmodulin-dependent serine/threonine kinase, or a fragment thereof, selected from the group consisting of:

5 (A) a polypeptide which is capable of inducing cell death (apoptosis) and comprises the amino acid sequence of SEQ ID NO:2;

(B) a polypeptide which has a property being capable of inducing cell death and has at least 85% sequence identity to the amino acid sequence of SEQ ID NO:2;

(C) a fragment of a polypeptide of SEQ ID NO:2 which is capable of inducing cell death;

10 (D) a fragment which is capable of inducing cell death and has at least 85% sequence identity to fragment (C);

(E) a fragment of a polypeptide of SEQ ID NO:2 which lacks the property of being capable of inducing cell death and which inhibits the ability of polypeptide (A) or (B) to induce cell death; and

15 (F) a fragment which lacks the property of being capable of inducing cell death and which inhibits the ability of polypeptide (A) or (B) to induce cell death, said fragment having at least 85% sequence identity to fragment (E).

2. An isolated DNA molecule comprising a nucleotide sequence encoding the polypeptide or fragment thereof according to claim 1.

20 3. The isolated DNA molecule according to claim 1, wherein said nucleotide sequence encodes the amino acid sequence of SEQ ID NO:2.

4. The isolated DNA molecule according to claim 3, wherein said nucleotide sequence corresponds to nucleotides 62 to 1141 of SEQ ID NO:1.

5. The isolated DNA molecule according to claim 3, which consists of the nucleotide sequence corresponding to nucleotides 62 to 1141 of SEQ ID NO:1.

6. An isolated DNA molecule which hybridizes to the DNA molecule of claim 5 under moderately stringent conditions and encodes a calmodulin-dependent serine/threonine kinase having the property of being capable of inducing cell death.

7. An isolated DNA molecule which hybridizes to the DNA molecule of claim 5 under highly stringent conditions and encodes a calmodulin-dependent serine/threonine kinase having the property of being capable of inducing cell death.

8. A polypeptide consisting of an amino acid sequence selected from the group consisting of amino acid residues 13 to 275 of SEQ ID NO:2 and an amino acid sequence having at least 85% sequence identity to residues 13 to 275 of SEQ ID NO:2.

9. An isolated DNA molecule comprising a nucleotide sequence encoding the polypeptide of claim 8.

10. The isolated DNA molecule according to claim 9, wherein said nucleotide sequence encodes the amino acid sequence corresponding to residues 13 to 275 of SEQ ID NO:2.

11. The isolated DNA molecule according to claim 10, wherein said nucleotide sequence is selected from the group consisting of nucleotides 98 to 886 of SEQ ID NO:1 and a nucleotide sequence which hybridizes to nucleotides 98 to 886 of SEQ ID NO:1 under moderately stringent conditions.

12. The isolated DNA molecule according to claim 11, wherein said nucleotide sequence hybridizes to nucleotides 98 to 886 of SEQ ID NO:1 under highly stringent conditions.

13. A polypeptide consisting of an amino acid sequence selected from the group consisting of amino acid residues 321 to 360 of SEQ ID NO:2 and an amino acid sequence having at least 85% sequence identity to residues 321 to 360 of SEQ ID NO:2.

14. An isolated DNA molecule comprising a nucleotide sequence encoding the
5 polypeptide of claim 13.

15. The isolated DNA molecule according to claim 14, wherein said nucleotide sequence encodes the amino acid sequence corresponding to residues 321 to 360 of SEQ ID NO:2.

16. The isolated DNA molecule according to claim 15, wherein said nucleotide
10 sequence is selected from the group consisting of nucleotides 1022 to 1141 of SEQ ID NO:1 and a nucleotide sequence which hybridizes to nucleotides 1022 to 1141 of SEQ ID NO:1 under moderately stringent conditions.

17. The isolated DNA molecule according to claim 16, wherein said nucleotide
15 sequence hybridizes to nucleotides 1022 to 1141 of SEQ ID NO:1 under highly stringent conditions.

18. A vector comprising the isolated DNA molecule according to any of claims 2-7, 9-12 and 14-17.

19. A host cell transformed with the isolated DNA molecule according to any of claims 2-7, 9-12, and 14-17.

20. A composition comprising a polypeptide according to any one of claims 1, 8 and 13, and a pharmaceutically acceptable excipient, carrier, diluent or auxiliary agent.

21. A molecule containing an antigen binding portion of an antibody which specifically recognizes the polypeptide according to any one of claims 1, 8 and 13 with the proviso that said antibody does not cross-react with DAP kinase or ZIP kinase.

22. The antibody according to claim 21, which is a monoclonal antibody.

23. A single stranded RNA molecule complementary to at least a portion of the isolated messenger RNA molecule which is the transcription product of the DNA sequence encoding a polypeptide of SEQ ID NO:2, wherein said complementary single stranded RNA molecule is capable of hybridizing to said isolated messenger RNA to prevent its translation
30 into said polypeptide of SEQ ID NO:2.

24. A method of neutralizing a messenger RNA molecule, which is the transcription product of the DNA sequence encoding a polypeptide of SEQ ID NO:2, comprising the step

of contacting the single stranded RNA molecule of claim 23 with the messenger RNA to neutralize the messenger RNA by hybridizing thereto and preventing its translation into the polypeptide of SEQ ID NO:2.

5 25. A method for screening individuals for a predisposition to cancer comprising the steps of:

(a) obtaining a sample of either genomic DNA from cells of the individual or cDNA produced from mRNA of said cells; and

(b) determining if there is a mutation in the nucleotide sequence of the DRP-1 gene.

10 26. The method according to claim 25 wherein a mutation in the nucleotide sequence of DRP-1 is determined by a process comprising the steps of:

(a) adding one or more nucleic acid probes to the sample of genomic DNA or cDNA, wherein each probe comprises a portion of the nucleotide sequence of DRP-1;

(b) providing conditions for hybridization between the nucleic acid probe or probes and the DNA of said samples; and

15 (c) determining on the basis of the hybridization whether there is a match between the sequence of the nucleic acid probe or probes and a sequence in the DNA of said sample, or whether there is a mismatch, a deletion or a mutation in the genomic DNA or cDNA and a predisposition to cancer of the tested individual.

FIG. 1

GACCGCGGCAGCTCAGCCTCCCGCCGATTGTATGTTCCAGGCCTCAATGAGGAGTCCAAA 60
M E P F K Q Q K V E D F Y D I G E E L G 20
CATGGAGCCATTCAAGCAGCAGAAGGTGGAGGACTTTTATGACATCGGAGAGGAGCTGGG 120
S G Q F A I V K K C R E K S T G L E Y A 40
GAGTGGCCAGTTTGCCATCGTGAAGAAGTGCCGGGAGAAGAGCACGGGGCTTGAGTATGC 180
A K F I K K R Q S R A S R R G V S R E E 60
AGCCAAGTTTCATCAAGAAGCGGCAGAGCCGGGCGAGCCGGCGCGGTGTGAGCCGGGAGGA 240
I E R E V S I L R Q V L H H N V I T L H 80
GATCGAGCGGGAGGTGAGCATCCTGCGGCAGGTGCTGCACCACAATGTCATCACGCTGCA 300
D V Y E N R T D V V L I L E L V S G G E 100
CGACGTCTATGAGAACCGCACCGACGTGGTGCTCATCCTTGAGCTAGTGTCTGGAGGAGA 360
L F D F L A Q K E S L S E E E A T S F I 120
GCTCTTCGATTTCTTGCGCCAGAAGGAGTCACTGAGTGAGGAGGAGGCCACCAGCTTCAT 420
K Q I L D G V N Y L H T K K I A H F D L 140
TAAGCAGATCCTGGATGGGGTGAACCTACCTTCACACAAAGAAAATTGCTCACTTTGATCT 480
K P E N I M L L D K N I P I P H I K L I 160
CAAGCCAGAAAACATTATGTTGTTAGACAAGAATATTCCCATTCCACACATCAAGCTGAT 540
D F G L A H E I E D G V E F K N I F G T 180
TGACTTTTGGTCTGGCTCACGAAATAGAAGATGGAGTTGAATTTAAGAATATTTTGGGAC 600
P E F V A P E I V N Y E P L G L E A D M 200
GCCGGAATTTGTTGCTCCAGAAATTGTGAACCTACGAGCCCCTGGGTCTGGAGGCTGACAT 660
W S I G V I T Y I L L S G A S P F L G D 220
GTGGAGCATAGGCGTCATCACCTACATCCTCTTAAGTGGAGCATCCCCTTTCCTGGGAGA 720
T K Q E T L A N I T S V S Y D F D E E F 240
CACGAAGCAGGAAACACTGGCAAATATCACATCAGTGAGTTACGACTTTGATGAGGAATT 780
F S H T S E L A K D F I R K L L V K E T 260
CTTCAGCCATACGAGCGAGCTGGCCAAGGACTTTATTCGGAAGCTTCTGGTTAAAGAGAC 840
R K R L T I Q E A L R H P W I T P V D N 280
CCGGAACGGCTCACAAATCCAAGAGGCTCTCAGACACCCCTGGATCACGCCGGTGGACAA 900
Q Q A M V R R E S V V N L E N F R K Q Y 300
CCAGCAAGCCATGGTGCAGGAGTCTGTGGTCAATCTGGAGAACTTCAGGAAGCAGTA 960
V R R R W K L S F S I V S L C N H L T R 320
TGTCGCGAGGCGGTGGAAGCTTTCCTTCAGCATCGTGTCCCTGTGCAACCACCTCACCCG 1020
S L M K K V H L R P D E D L R N C E S D 340
CTCGCTGATGAAGAAGGTGCACCTGAGGCCGGATGAGGACCTGAGGAACTGTGAGAGTGA 1080
T E E D I A R R K A L H P R R R S S T S 360
CACTGAGGAGGACATcGCCAGGAGGAAAGCCCTCCACCCACGGAGGAGGAGCAGCACCTC 1140
CTAAcTGGCCTGACCTGCAGTGGCCGCCAGGGAGGTTTGGGCCCAGCGGGGCTCCCTTCT 1200
GTGCAGACTTTTGGACCCAGCTCAGCACCCAGCACCCGGGCGTCCTGAGCACTTTGCAAGA 1260
GAGATGGGCCCCAAGGAATTCAGAAGAGCTTGCAAGCAAGCCAGGAGACCCTGGGAGCTGT 1320
GGCTGTcTTCTGTGGAGGAGGCTCCAGCATTCCCAAAGCTCTTAATTCTCCATAAAATGG 1380
GCTTTCCTCTGTCTGCCATCCTCAGAGTCTGGGGTGGGAGTGTGGAcTTAGGAAAACAAT 1440
ATAAAGGACATCCTCATCATCACGGGGTGAAGGTCAGAGTAAGGCAGCCTTCTTCACAGG 1500
CTGAGGGGGTTTCAAGAACAGCCTGGCCAAAAATTACACCAGAGAGACAGAGTCCTCCCCA 1560
TTGGGAACAGGGTGATTGAGGAAAGTGAACCTTgGGTGTGAGGGACCAATCCTGTGACCT 1620
CCCAGAACCATGGAAGCCAGGACGTCAGGCTGaCCAACACCTCAGACCTTCTGAAGCAGC 1680
CCATTGcTGGCCCGCCATGTTGTAATTTTGCTCATTTTTATTAAActtctggtttacctg 1740
aa 1742

2 / 16

FIG. 2A

DAP-kinase	13	YDTGEEELGSGQFAVVKKCREKSTGLQYAAKFIK
ZIP-kinase	13	YEMGEEELGSGQFAIVRKCRQKGTGKEYAAKFIK
DRP-1	13	YDIGEEELGSGQFAIVKKCREKSTGLE YAAKFIK
DRAK1	61	LCPGRELGRGKFAVVVKCIKKDSGKEFAAKFMR
DRAK2	33	ILTSKELGRGKFAVVVRQCI SKSTGQ EYAAKFLK
DAP-kinase	46	KRR TKSSRRGVSRREDIEREVSILKEI-QHPNVI
ZIP-kinase	46	KRRLSSSRGVSRREEIEREVNILEI-RHPNII
DRP-1	46	KRQSRASRRGVSRREEIEREVSILRQV-LHHNVI
DRAK1	94	KRR----KGQDCRMEI IHEIAVLELAQDNPWVI
DRAK2	66	KRR----RGQDCRAEILHEIAVLELAKSCPRVI
DAP-kinase	78	TLHEVYENKTDVILILELVAGGELFD-FLAEK-
ZIP-kinase	78	TLHDIENKTDVVLILELVSGGELFD-FLAEK-
DRP-1	78	TLHDVYENRTDVVLILELVSGGELFD-FLAQK-
DRAK1	123	NLHEVYETASEMILVLEYAAGGEIFDQCVADRE
DRAK2	95	NLHEVYENTSEIILILEY AAGGEIFSLCLPELA
DAP-kinase	109	ESLTEEEATEFLKQILNGVYYLHSLQIAHF DLK
ZIP-kinase	109	ESLTEDEATQFLKQILDGVHYLH SKRIAHF DLK
DRP-1	109	ESLSEEEATSFIKQILDGVN YLHTKKIAHF DLK
DRAK1	156	EAFKEKDVQRLMRQILEGVHFLHTRD VVHLDL K
DRAK2	128	EMVSENDVIRLIKQILEGVYYLH QNNIVHLDL K
DAP-kinase	142	PENIMLLDRNVPKPRIKIIDFG LAHKIDFGNEF
ZIP-kinase	142	PENIMLLDKNVPNPRIKLIDFG IAHKIEAGNEF
DRP-1	142	PENIMLLDKNIPIPHIKLIDFG LAHEIEDGV EF
DRAK1	189	PQNILLTSESPLGD-IKIVDFGLSRILKNSEEL
DRAK2	161	PQNILLSSIYPLGD-IKIVDFGMSRKIGHACEL
DAP-kinase	175	KNIFGTPEFVAPETVNYEPLGLEADMWSIGVIT
ZIP-kinase	175	KNIFGTPEFVAPETVNYEPLGLEADMWSIGVIT
DRP-1	175	KNIFGTPEFVAPETVNYEPLGLEADMWSIGVIT
DRAK1	221	REIMGTPEYVAPETLSYDPISMATDMWSIGVLT
DRAK2	193	REIMGTPEYLAPEILNYDPITTATDMWNIGITIA
DAP-kinase	208	YILLSGASPFLLGDTKQETLANV SAVNYEFED EY
ZIP-kinase	208	YILLSGASPFLLGETKQETLTNIS SAVNYDFDEEY
DRP-1	208	YILLSGASPFLLGDTKQETLANITSVSYDFDEEF
DRAK1	254	YVMLTGISPFLLGNDKQETFLNISQMNLSYSEEE
DRAK2	226	YMLLTHTSPFVGEDNQETYLNISQVNV DYSEET
DAP-kinase	241	FSNTSALAKDFIRRLLVKDPKKRMTIQDSLOHP
ZIP-kinase	241	FSNTSELAKDFIRRLLVKDPKKRMTIAQSLEHS
DRP-1	241	FSHTSELAKDFIRKLLVKETRKRLTIQEALRHP
DRAK1	287	FDVLSESAVDFIRTLLVKKPEDRATAEELKHP
DRAK2	259	FSSVSQLATDFIQSLLVKNPEKRPTAEICLSHS
DAP-kinase	274	WI
ZIP-kinase	274	WI
DRP-1	274	WI
DRAK1	320	WL
DRAK2	292	WL

3 / 16


FIG. 2B

DAP-kinase	1	NMEKFKK---FAARKKWKQSVRLISLCORLSR	29
DRP-1	1	NLENFRK---QYVRRRWKLSFSIVSLCNHLTR	29
smMLCK	1	SKDRMKK---YMARRKWQKTGHAVRAICRLSS	29
CaMKIIa	1	TVDCLKK---LNARRKLGAILTTMLATRNFS	29
CaMKI	1	VSEQIKK---NFAKSKWKQAFNAT-AVVRHMR	28
CaMKIV	1	MDTAQKKLQEFNARRKKA AVKAVVASSRLGS	32
ZIP-kinase	1	GEDSGRK---PERRRLKTTTRLKEYTIKSHSS	28

FIG. 3C

FIG. 3B

FIG. 3A



5 / 16

FIG. 4A


MOCK

*FIG. 4B*

DRP-1

*FIG. 4C*

DAP-kinase		DRP-1	
InSol	Sol	InSol	Sol






FIG. 5A

FIG. 5B




FIG. 6B

FIG. 6D





FIG. 6A

FIG. 6C

FIG. 7



9 / 16





FIG. 8A*FIG. 8B*

FIG. 9A



11 / 16

FIG. 9B

12 / 16

FIG. 10A




FIG. 10B





FIG. 11A



14 / 16


FIG. 11B

FIG. 12A





FIG. 12B



SEQUENCE LISTING

<110> KIMCHI, Adi
 MCINNIS A., Patricia
 YEDA RESEARCH AND DEVELOPMENT COMPANY LTD.

<120> DAP-KINASE RELATED PROTEIN

<130> KIMCHI2A

<140> 00

<141> 1999-06-15

<150> 60/089,294

<151> 1998-06-15

<160> 14

<170> PatentIn Ver. 2.0

<210> 1

<211> 1742

<212> DNA

<213> Human

<220>

<221> CDS

<222> (62)..(1141)

<400> 1

gaccgcggca gctcagcctc ccgccgattg tatgttccag gcctcaatga ggagtccaaa 60

c atg gag cca ttc aag cag cag aag gtg gag gac ttt tat gac atc gga 109

Met Glu Pro Phe Lys Gln Gln Lys Val Glu Asp Phe Tyr Asp Ile Gly

1 5 10 15

gag gag ctg ggg agt ggc cag ttt gcc atc gtg aag aag tgc cgg gag 157

Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Lys Lys Cys Arg Glu

20 25 30

aag agc acg ggg ctt gag tat gca gcc aag ttc atc aag aag cgg cag 205

Lys Ser Thr Gly Leu Glu Tyr Ala Ala Lys Phe Ile Lys Lys Arg Gln

35 40 45

agc cgg gcg agc cgg cgc ggt gtg agc cgg gag gag atc gag cgg gag 253

Ser Arg Ala Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu Arg Glu

50 55 60

gtg agc atc ctg cgg cag gtg ctg cac cac aat gtc atc acg ctg cac	301
Val Ser Ile Leu Arg Gln Val Leu His His Asn Val Ile Thr Leu His	
65 70 75 80	
 gac gtc tat gag aac cgc acc gac gtg gtg cac atc ctt gag cta gtg	349
Asp Val Tyr Glu Asn Arg Thr Asp Val Val His Ile Leu Glu Leu Val	
85 90 95	
 tct gga gga gag ctc ttc gat ttc ctg gcc cag aag gag tca ctg agt	397
Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Gln Lys Glu Ser Leu Ser	
100 105 110	
 gag gag gag gcc acc agc ttc att aag cag atc ctg gat ggg gtg aac	445
Glu Glu Glu Ala Thr Ser Phe Ile Lys Gln Ile Leu Asp Gly Val Asn	
115 120 125	
 tac ctt cac aca aag aaa att gct cac ttt gat ctc aag cca gaa aac	493
Tyr Leu His Thr Lys Lys Ile Ala His Phe Asp Leu Lys Pro Glu Asn	
130 135 140	
 att atg ttg tta gac aag aat att ccc att cca cac atc aag ctg att	541
Ile Met Leu Leu Asp Lys Asn Ile Pro Ile Pro His Ile Lys Leu Ile	
145 150 155 160	
 gac ttt ggt ctg gct cac gaa ata gaa gat gga gtt gaa ttt aag aat	589
Asp Phe Gly Leu Ala His Glu Ile Glu Asp Gly Val Glu Phe Lys Asn	
165 170 175	
 att ttt ggg acg ccg gaa ttt gtt gct cca gaa att gtg aac tac gag	637
Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn Tyr Glu	
180 185 190	
 ccc ctg ggt ctg gag gct gac atg tgg agc ata ggc gtc atc acc tac	685
Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly Val Ile Thr Tyr	
195 200 205	
 atc ctc tta agt gga gca tcc cct ttc ctg gga gac acg aag cag gaa	733
Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Asp Thr Lys Gln Glu	
210 215 220	
 aca ctg gca aat atc aca tca gtg agt tac gac ttt gat gag gaa ttc	781
Thr Leu Ala Asn Ile Thr Ser Val Ser Tyr Asp Phe Asp Glu Glu Phe	
225 230 235 240	
 ttc agc cat acg agc gag ctg gcc aag gac ttt att cgg aag ctt ctg	829
Phe Ser His Thr Ser Glu Leu Ala Lys Asp Phe Ile Arg Lys Leu Leu	
245 250 255	

gtt aaa gag acc cgg aaa cgg ctc aca atc caa gag gct ctc aga cac 877
 Val Lys Glu Thr Arg Lys Arg Leu Thr Ile Gln Glu Ala Leu Arg His
 260 265 270

ccc tgg atc acg ccg gtg gac aac cag caa gcc atg gtg cga cgg gag 925
 Pro Trp Ile Thr Pro Val Asp Asn Gln Gln Ala Met Val Arg Arg Glu
 275 280 285

tct gtg gtc aat ctg gag aac ttc agg aag cag tat gtc cgc agg cgg 973
 Ser Val Val Asn Leu Glu Asn Phe Arg Lys Gln Tyr Val Arg Arg Arg
 290 295 300

tgg aag ctt tcc ttc agc atc gtg tcc ctg tgc aac cac ctc acc cgc 1021
 Trp Lys Leu Ser Phe Ser Ile Val Ser Leu Cys Asn His Leu Thr Arg
 305 310 315 320

tcg ctg atg aag aag gtg cac ctg agg ccg gat gag gac ctg agg aac 1069
 Ser Leu Met Lys Lys Val His Leu Arg Pro Asp Glu Asp Leu Arg Asn
 325 330 335

tgt gag agt gac act gag gag gac atc gcc agg agg aaa gcc ctc cac 1117
 Cys Glu Ser Asp Thr Glu Glu Asp Ile Ala Arg Arg Lys Ala Leu His
 340 345 350

cca cgg agg agg agc agc acc tcc taactggcct gacctgcagt ggccgccagg 1171
 Pro Arg Arg Arg Ser Ser Thr Ser
 355 360

gaggtttggg cccagcgggg ctcccttctg tgcagacttt tggacccagc tcagcaccag 1231

caccggggcg tcctgagcac tttgcaagag agatggggccc aaggaattca gaagagcttg 1291

caggcaagcc aggagaccct gggagctgtg gctgtcttct gtggaggagg ctccagcatt 1351

cccaaagetc ttaattctcc ataaaatggg ctttctctg tctgccatcc tcagagtctg 1411

ggggtgggagt gtggacttag gaaaacaata taaaggacat cctcatcatc acgggggtgaa 1471

ggtcagagta aggcagcctt cttcacaggc tgaggggggtt cagaaccagc ctggccaaaa 1531

attacaccag agagacagag tcctcccat tgggaacagg gtgattgagg aaagtgaacc 1591

ttgggtgtga gggaccaatc ctgtgacctc ccagaaccat ggaagccagg acgtcaggct 1651

gaccaacacc tcagaccttc tgaagcagcc cattgctggc ccgccatgtt gtaattttgc 1711

tcatttttat taaacttctg gtttacctga a 1742

<210> 2

<211> 360

<212> PRT

<213> Human

<400> 2

Met Glu Pro Phe Lys Gln Gln Lys Val Glu Asp Phe Tyr Asp Ile Gly

1

5

10

15

Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Lys Lys Cys Arg Glu

20

25

30

Lys Ser Thr Gly Leu Glu Tyr Ala Ala Lys Phe Ile Lys Lys Arg Gln

35

40

45

Ser Arg Ala Ser Arg Arg Gly Val Ser Arg Glu Glu Ile Glu Arg Glu

50

55

60

Val Ser Ile Leu Arg Gln Val Leu His His Asn Val Ile Thr Leu His

65

70

75

80

Asp Val Tyr Glu Asn Arg Thr Asp Val Val His Ile Leu Glu Leu Val

85

90

95

Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Gln Lys Glu Ser Leu Ser

100

105

110

Glu Glu Glu Ala Thr Ser Phe Ile Lys Gln Ile Leu Asp Gly Val Asn

115

120

125

Tyr Leu His Thr Lys Lys Ile Ala His Phe Asp Leu Lys Pro Glu Asn

130

135

140

Ile Met Leu Leu Asp Lys Asn Ile Pro Ile Pro His Ile Lys Leu Ile

145

150

155

160

Asp Phe Gly Leu Ala His Glu Ile Glu Asp Gly Val Glu Phe Lys Asn

165

170

175

Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn Tyr Glu

180

185

190

Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly Val Ile Thr Tyr

195

200

205

Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Asp Thr Lys Gln Glu

210

215

220

Thr Leu Ala Asn Ile Thr Ser Val Ser Tyr Asp Phe Asp Glu Glu Phe
 225 230 235 240

Phe Ser His Thr Ser Glu Leu Ala Lys Asp Phe Ile Arg Lys Leu Leu
 245 250 255

Val Lys Glu Thr Arg Lys Arg Leu Thr Ile Gln Glu Ala Leu Arg His
 260 265 270

Pro Trp Ile Thr Pro Val Asp Asn Gln Gln Ala Met Val Arg Arg Glu
 275 280 285

Ser Val Val Asn Leu Glu Asn Phe Arg Lys Gln Tyr Val Arg Arg Arg
 290 295 300

Trp Lys Leu Ser Phe Ser Ile Val Ser Leu Cys Asn His Leu Thr Arg
 305 310 315 320

Ser Leu Met Lys Lys Val His Leu Arg Pro Asp Glu Asp Leu Arg Asn
 325 330 335

Cys Glu Ser Asp Thr Glu Glu Asp Ile Ala Arg Arg Lys Ala Leu His
 340 345 350

Pro Arg Arg Arg Ser Ser Thr Ser
 355 360

<210> 3

<211> 263

<212> PRT

<213> Human

<400> 3

Tyr Asp Thr Gly Glu Glu Leu Gly Ser Gly Gln Phe Ala Val Val Lys
 1 5 10 15

Lys Cys Arg Glu Lys Ser Thr Gly Leu Gln Tyr Ala Ala Lys Phe Ile
 20 25 30

Lys Lys Arg Arg Thr Lys Ser Ser Arg Arg Gly Val Ser Arg Glu Asp
 35 40 45

Ile Glu Arg Glu Val Ser Ile Leu Lys Glu Ile Gln His Pro Asn Val
 50 55 60

Ile Thr Leu His Glu Val Tyr Glu Asn Lys Thr Asp Val Ile Leu Ile

65 70 75 80
 Leu Glu Leu Val Ala Gly Gly Glu Leu Phe Asp Phe Leu Ala Glu Lys
 85 90 95
 Glu Ser Leu Thr Glu Glu Glu Ala Thr Glu Phe Leu Lys Gln Ile Leu
 100 105 110
 Asn Gly Val Tyr Tyr Leu His Ser Leu Gln Ile Ala His Phe Asp Leu
 115 120 125
 Lys Pro Glu Asn Ile Met Leu Leu Asp Arg Asn Val Pro Lys Pro Arg
 130 135 140
 Ile Lys Ile Ile Asp Phe Gly Leu Ala His Lys Ile Asp Phe Gly Asn
 145 150 155 160
 Glu Phe Lys Asn Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile
 165 170 175
 Val Asn Tyr Glu Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly
 180 185 190
 Val Ile Thr Tyr Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Asp
 195 200 205
 Thr Lys Gln Glu Thr Leu Ala Asn Val Ser Ala Val Asn Tyr Glu Phe
 210 215 220
 Glu Asp Glu Tyr Phe Ser Asn Thr Ser Ala Leu Ala Lys Asp Phe Ile
 225 230 235 240
 Arg Arg Leu Leu Val Lys Asp Pro Lys Lys Arg Met Thr Ile Gln Asp
 245 250 255
 Ser Leu Gln His Pro Trp Ile
 260

<210> 4
 <211> 263
 <212> PRT
 <213> Human

<400> 4
 Tyr Glu Met Gly Glu Glu Leu Gly Ser Gly Gln Phe Ala Ile Val Arg
 1 5 10 15

Lys Cys Arg Gln Lys Gly Thr Gly Lys Glu Tyr Ala Ala Lys Phe Ile
 20 25 30

Lys Lys Arg Arg Leu Ser Ser Ser Arg Arg Gly Val Ser Arg Glu Glu
 35 40 45

Ile Glu Arg Glu Val Asn Ile Leu Arg Glu Ile Arg His Pro Asn Ile
 50 55 60

Ile Thr Leu His Asp Ile Phe Glu Asn Lys Thr Asp Val Val Leu Ile
 65 70 75 80

Leu Glu Leu Val Ser Gly Gly Glu Leu Phe Asp Phe Leu Ala Glu Lys
 85 90 95

Glu Ser Leu Thr Glu Asp Glu Ala Thr Gln Phe Leu Lys Gln Ile Leu
 100 105 110

Asp Gly Val His Tyr Leu His Ser Lys Arg Ile Ala His Phe Asp Leu
 115 120 125

Lys Pro Glu Asn Ile Met Leu Leu Asn Lys Asn Val Pro Asn Pro Arg
 130 135 140

Ile Lys Leu Ile Asp Phe Gly Ile Ala His Lys Ile Glu Ala Gly Asn
 145 150 155 160

Glu Phe Lys Asn Ile Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Ile
 165 170 175

Val Asn Tyr Glu Pro Leu Gly Leu Glu Ala Asp Met Trp Ser Ile Gly
 180 185 190

Val Ile Thr Tyr Ile Leu Leu Ser Gly Ala Ser Pro Phe Leu Gly Glu
 195 200 205

Thr Lys Gln Glu Thr Leu Thr Asn Ile Ser Ala Val Asn Tyr Asp Phe
 210 215 220

Asp Glu Glu Tyr Phe Ser Asn Thr Ser Glu Leu Ala Lys Asp Phe Ile
 225 230 235 240

Arg Arg Leu Leu Val Lys Asp Pro Lys Arg Arg Met Thr Ile Ala Gln
 245 250 255

Ser Leu Glu His Ser Trp Ile
 260

<210> 5

<211> 261

<212> PRT

<213> Human

<400> 5

Leu Cys Pro Gly Arg Glu Leu Gly Arg Gly Lys Phe Ala Val Val Arg
 1 5 10 15

Lys Cys Ile Lys Lys Asp Ser Gly Lys Glu Phe Ala Ala Lys Phe Met
 20 25 30

Arg Lys Arg Arg Lys Gly Gln Asp Cys Arg Met Glu Ile Ile His Glu
 35 40 45

Ile Ala Val Leu Glu Leu Ala Gln Asp Asn Pro Trp Val Ile Asn Leu
 50 55 60

His Glu Val Tyr Glu Thr Ala Ser Glu Met Ile Leu Val Leu Glu Tyr
 65 70 75 80

Ala Ala Gly Gly Glu Ile Phe Asp Gln Cys Val Ala Asp Arg Glu Glu
 85 90 95

Ala Phe Lys Glu Lys Asp Val Gln Arg Leu Met Arg Gln Ile Leu Glu
 100 105 110

Gly Val His Phe Leu His Thr Arg Asp Val Val His Leu Asp Leu Lys
 115 120 125

Pro Gln Asn Ile Leu Leu Thr Ser Glu Ser Pro Leu Gly Asp Ile Lys
 130 135 140

Ile Val Asp Phe Gly Leu Ser Arg Ile Leu Lys Asn Ser Glu Glu Leu
 145 150 155 160

Arg Glu Ile Met Gly Thr Pro Glu Tyr Val Ala Pro Glu Ile Leu Ser
 165 170 175

Tyr Asp Pro Ile Ser Met Ala Thr Asp Met Trp Ser Ile Gly Val Leu
 180 185 190

Thr Tyr Val Met Leu Thr Gly Ile Ser Pro Phe Leu Gly Asn Asp Lys
 195 200 205

Gln Glu Thr Phe Leu Asn Ile Ser Gln Met Asn Leu Ser Tyr Ser Glu
 210 215 220

Glu Glu Phe Asp Val Leu Ser Glu Ser Ala Val Asp Phe Ile Arg Thr
 225 230 235 240

Leu Leu Val Lys Lys Pro Glu Asp Arg Ala Thr Ala Glu Glu Cys Leu
 245 250 255

Lys His Pro Trp Leu
 260

<210> 6

<211> 261

<212> PRT

<213> Human

<400> 6

Ile Leu Thr Ser Lys Glu Leu Gly Arg Gly Lys Phe Ala Val Val Arg
 1 5 10 15

Gln Cys Ile Ser Lys Ser Thr Gly Gln Glu Tyr Ala Ala Lys Phe Leu
 20 25 30

Lys Lys Arg Arg Arg Gly Gln Asp Cys Arg Ala Glu Ile Leu His Glu
 35 40 45

Ile Ala Val Leu Glu Leu Ala Lys Ser Cys Pro Arg Val Ile Asn Leu
 50 55 60

His Glu Val Tyr Glu Asn Thr Ser Glu Ile Ile Leu Ile Leu Glu Tyr
 65 70 75 80

Ala Ala Gly Gly Glu Ile Phe Ser Leu Cys Leu Pro Glu Leu Ala Glu
 85 90 95

Met Val Ser Glu Asn Asp Val Ile Arg Leu Ile Lys Gln Ile Leu Glu
 100 105 110

Gly Val Tyr Tyr Leu His Gln Asn Asn Ile Val His Leu Asp Leu Lys
 115 120 125

Pro Gln Asn Ile Leu Leu Ser Ser Ile Tyr Pro Leu Gly Asp Ile Lys
 130 135 140

Ile Val Asp Phe Gly Met Ser Arg Lys Ile Gly His Ala Cys Glu Leu
 145 150 155 160

Arg Glu Ile Met Gly Thr Pro Glu Tyr Leu Ala Pro Glu Ile Leu Asn

165 170 175
 Tyr Asp Pro Ile Thr Thr Ala Thr Asp Met Trp Asn Ile Gly Ile Ile
 180 185 190
 Ala Tyr Met Leu Leu Thr His Thr Ser Pro Phe Val Gly Glu Asp Asn
 195 200 205
 Gln Glu Thr Tyr Leu Asn Ile Ser Gln Val Asn Val Asp Tyr Ser Glu
 210 215 220
 Glu Thr Phe Ser Ser Val Ser Gln Leu Ala Thr Asp Phe Ile Gln Ser
 225 230 235 240
 Leu Leu Val Lys Asn Pro Glu Lys Arg Pro Thr Ala Glu Ile Cys Leu
 245 250 255
 Ser His Ser Trp Leu
 260

<210> 7
 <211> 29
 <212> PRT
 <213> Human

<400> 7
 Asn Met Glu Lys Phe Lys Lys Phe Ala Ala Arg Lys Lys Trp Lys Gln
 1 5 10 15
 Ser Val Arg Leu Ile Ser Leu Cys Gln Arg Leu Ser Arg
 20 25

<210> 8
 <211> 29
 <212> PRT
 <213> Human

<400> 8
 Asn Leu Glu Asn Phe Arg Lys Gln Tyr Val Arg Arg Arg Trp Lys Leu
 1 5 10 15
 Ser Phe Ser Ile Val Ser Leu Cys Asn His Leu Thr Arg
 20 25

<210> 9

<211> 29
<212> PRT
<213> Human

<400> 9
Thr Cys Asp Cys Leu Lys Lys Leu Asn Ala Arg Arg Lys Leu Lys Gly
1 5 10 15

Ala Ile Leu Thr Thr Met Leu Ala Thr Arg Asn Phe Ser
20 25

<210> 10
<211> 28
<212> PRT
<213> Human

<400> 10
Val Ser Glu Gln Ile Lys Lys Asn Phe Ala Lys Ser Lys Trp Lys Gln
1 5 10 15

Ala Phe Asn Ala Thr Ala Val Val Arg His Met Arg
20 25

<210> 11
<211> 32
<212> PRT
<213> Human

<400> 11
Met Asp Thr Ala Gln Lys Lys Leu Gln Glu Phe Asn Ala Arg Arg Lys
1 5 10 15

Leu Lys Ala Ala Val Lys Ala Val Val Ala Ser Ser Arg Leu Gly Ser
20 25 30

<210> 12
<211> 28
<212> PRT
<213> Human

<400> 12
Gly Glu Asp Ser Gly Arg Lys Pro Glu Arg Arg Arg Leu Lys Thr Thr

1

5

10

15

Arg Leu Lys Glu Tyr Thr Ile Lys Ser His Ser Ser
20 25

<210> 13

<211> 20

<212> DNA

<213> Human

<400> 13

ggccggatga ggacctgagg

20

<210> 14

<211> 21

<212> DNA

<213> Human

<400> 14

tccacatccc accccagact c

21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 9/12, 1/20, 5/00, 15/00, C12Q 1/68; C07H 21/04; A61K 38/51

US CL : 435/194, 320.2, 325, 252.3, 6; 424/94.5; 536/23.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/194, 320.2, 325, 252.3, 6; 424/94.5; 536/23.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9510630 A (YEDA RESEARCH AND DEVELOPMENT CO. LTD.) 20 April 1995 (20.04.95), see the entire article specially compare their kinase domain (AA 13-275) to a fragment of DRP-1 and its homologs in claims 1C-D.	1-7 and 20
X	DEISS et al. Identification of a novel serine/threonine kinase and a novel 15-kD protein as potential mediators of the γ interferon-induced cell death. Genes & Development. 01 January 1995, Vol. 9, pages 15-30, see figures 4 and 7.	8-12, 20

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 NOVEMBER 1999

Date of mailing of the international search report

19 NOV 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MARYAM MONSHIPOURI CV

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database EST on GenCore version 4.5, Accession No. H27327, HILLIER et al. 'The WashU-Merck Project', definition: y116d12.r1 Soars breast 2NbHBst Homosapiens cDNA clone. 12 July 1995, see their residues 81-259 for 100% identity to residues 1300-1478 of SEQ ID NO:1.	23
---		----
Y		24

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13411

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-20, 23-24

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/13411

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Detailed Reasons For Holding Lack of Unity of Invention:

This International Preliminary Examination Authority has found 3 inventions claimed in the International Application covered by the claims indicated below:

Group I, claims 1-20 and 23-24, drawn to isolated DNA molecules encoding calmodulin-dependent serine/threonine kinases (DRP-1) and their fragments, vectors and host cells comprising said DNA molecules, single stranded antisense RNA molecules complementary to at least a portion of the transcription product of said DNA molecules, methods of neutralizing a messenger RNA molecule, which is the transcription product of DRP-1 encoding gene comprising contacting said antisense RNA molecules to messenger RNA molecules in order to prevent translation, calmodulin-dependent serine/threonine (DRP-1) kinases and their fragments together with compositions comprising said kinases and their fragments.

Group II, claims 21-22, drawn to a molecule containing an antigen binding portion of an antibody which specifically binds said DRP-1 kinases.

Group III, claims 25-26, drawn to a method of screening cancer patients comprising obtaining a sample of either genomic DNA or cDNA of cancer cells and determining if a mutation occurred in the gene encoding DRP-1.

The inventions listed as Groups I-III do not relate to a single inventive concept because they are considered to be three categories of invention and are not drawn to combination of categories (i.e. categories 1-5), specified in 37 CFR section 1.475(B).

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.